



CHEMISTRY

2. Analytical Biogeochemistry



Baffin Island Oil Spill Project

WORKING REPORT SERIES
83-2

1983 STUDY RESULTS

BAFFIN ISLAND OIL SPILL PROJECT
WORKING REPORT SERIES

The Baffin Island Oil Spill (BIOS) Project is a multidisciplinary program of research on arctic marine oilspill fate, effects and countermeasures. The Project commenced in the spring of 1980 and has now completed the fourth and final year of planned field work at an experimental site located on the northern end of Baffin Island, Canada. The results of work performed in each of the various study components under the Project have been made available on a yearly basis through this working report series. This has been done prior to a complete integration of findings and interpretation with respect to the Project objectives. The working report series should therefore be considered as Interim or data reports. The contents do not necessarily reflect the views Or policies of the BIOS Project management or funders.

Reprint or republication of contents should not be made without the consent of the author or the BIOS Project.

For further information on the BIOS Project write the BIOS Project Manager, c/o Environmental Emergencies Technology Division, Environmental Protection Service, Ottawa, Ontario, Canada K1A 1C8.

The Baffin Island Oil Spill Project has been funded and supported,

In Canada by:

Department of the Environment (Environmental Protection Service)
Canadian Offshore Oil Spill Research Association
Department of Indian and Northern Affairs
Canadian Coast Guard
Environmental Studies Revolving Fund (IAND)
Environmental Studies Revolving Fund (EMR)
Petro Canada Resources
Department of Fisheries and Oceans
Offshore Labrador Biological Study
Texaco Canada
Polar Continental Shelf
Pond Inlet Hamlet Council

In the United States by:

National Oceanographic and Atmospheric Administration (USA)
United States Coast Guard Office of Research and Development
American Petroleum Institute
Exxon

In the United Kingdom by:

BP International (London)

In Norway by:

Royal Norwegian Ministry of Environment (FOH and PFO)

CORRECT CITATION FOR THIS PUBLICATION:

BOEHM, P., W. STEINHAUER, D. COBB, S. DUFFY AND J. BROWN. 1984.
CHEMISTRY 2: ANALYTICAL BIOGEOCHEMISTRY -1983 STUDY RESULTS.
BAFFIN ISLAND OIL SPILL WORKING REPORT 83-2; 139p. ENVIRONMENTAL
PROTECTION SERVICE, ENVIRONMENT CANADA, OTTAWA.

BAFFIN ISLAND OIL SPILL PROJECT
CHEMISTRY COMPONENT-A
ANALYTICAL BIOGEOCHEMISTRY
REPORT ON 1983 FIELD EXPERIMENTS

FINAL REPORT
CONTRACT NO. 0S283-00039

PREPARED FOR

BIOS PROJECT OFFICE
ENVIRONMENT PROTECTION SERVICE .
804, 9942108 Street
Edmonton, Alberta T5K 235
Canada
Attn: Mr. Gary Sergy

PREPARED BY:

Paul D. Boehm, **William Steinhauer**, Donald Cobb,
Suzanne **Duffy** and John Brown

BATTELLE
New England Marine Research Laboratory
397 Washington Street
Duxbury, MA 02332

May 31, 1984

Battelle is not engaged **in** research for advertising, sales promotion, or **publicity** purposes, and this report may not be **reproduced** in full or in part for such **purposes.**

TABLE OF CONTENTS

| | <u>Page</u> |
|--|-------------|
| EXECUTIVE SUMMARY . . . *.0..... | i |
| SECTION ONE INTRODUCTION0.0 | 1 |
| 1.1 Project Goals | 1 |
| 1.2 Technical Plan | 1 |
| SECTION TWO SAMPLING AND ANALYTICAL METHODS | 4 |
| 2.1 Sampling** | 4 |
| 2.1.1 Seawater Sampling | 4 |
| 2.1.2 Sediment Sampling | 8 |
| 2.1.3 Benthic Animal Sampling0.... 0.... | 9 |
| 2.2 Analytical Methods* | 9 |
| 2.2.1 Sample Processing0... . . . 0.... | 11 |
| 2.2.1.1 Water Samples (16 Liter) . . . *.... . . 00... | 11 |
| 2.2.1.2 Large Volume Water Samples0 . . * . . | 13 |
| 2.2.1.3 Sediment Sample Processing*.. | 13 |
| 2.2.1.3a Surface Sediments (0-2 cm) | 13 |
| 2.2.1.3b Surface Floe Samples . . . 0.... | 13 |
| 2.2.1.3c Oiled Beach Sediments | 15 |
| 2.2.1.4 Benthic Animal Tissue Processing* | 15 |
| 2.2.2 Sample Analysis . . * 00 * | 16 |
| 2.2.2.1 UV/F Analysis . . 0....0 . . 0.... * | 16 |
| 2.2.2.2 Fractionation0 . . 00.... . . * . . * | 16 |
| 2.2.2.3 Gas Chromatography (GC ²) Analysis . . . *** | 18 |
| 2.2.2.4 Gas Chromatography/Mass Spectrometry (GC ² /MS) | 22 |
| SECTION THREE RESULTS (NEARSHORE STUDY)0....0 . . . * | 25 |
| 3.1 Water Column** . . . 0...0.. | 25 |
| 3.1.1 Oil on the Water's Surface (Surface Slick)0. | 25 |

TABLE OF CONTENTS
(Continued)

| | <u>Page</u> |
|--|-------------|
| 3.1.2 Oil in the Water Column* | 30 |
| 3.1.2.1 16 Liter Samples 0....000.. | 30 |
| 3.1.2.2 Large-Volume Samples | 32 |
| 3.2 Oil in the Sediments**..... | 35 |
| 3.2.1 Bay 110..... | 35 |
| 3.2.1.1 Tissue Plots | 35 |
| 3.2.1.1a Oil Concentrations by UV/F | 35 |
| 3.2.1.1b Oil Composition by GC² | 38 |
| 3.2.1.1c Aromatic Hydrocarbon Composition by GC²/MS | 38 |
| 3.2.1.2 Surface Floe Samples* | 38 |
| 3.2.1.2a Oil Concentration by UV/F0.....0.0. | 38 |
| 3.2.1.2b Oil Composition by GC² | 42 |
| 3.2. 1.2c Aromatic Hydrocarbon Composition by GC²/MS | 42 |
| 3.2.1.3 Benthic Transects0..... | 42 |
| 3.2. 1.3a Oil Concentration by UV/F0..0. | 42 |
| 3.2.1.3b Oil Composition by GC² | 45 |
| 3.2. 1.3c Aromatic Hydrocarbon Composition BY GC²/MS | 45 |
| 3.2.1.4 Microbiology Transects0.....* | 45 |
| 3.2. 1.4a Oil Concentration and Composition by GC² | 45 |
| 3.2.1.5 Sediment Cores | 49 |
| 3.2.1.5a Oil Composition by GC²* | 49 |
| 3.2.1.5b Aromatic Hydrocarbon Composition by GC²/MS | 53 |
| 3.2.1.6 Deep Sediments (35 m)0..... .. | 53 |
| 3.2.1.6a Oil Concentrations by UV/F | 53 |
| 3.2.1.6b Oil Composition by GC² | 53 |
| 3.2.1.6c Aromatic Hydrocarbon Composition by GC²/MS | 61 |
| 3.2.1.7 Bay 11 Beach Sediments000 | 61 |

TABLE OF CONTENTS
(Continued)

| | <u>Page</u> |
|--|-------------|
| 3.2.2 Bay9 | 61 |
| 3.2.2.1 Tissue Plots* | 64 |
| 3.2.2.1a Oil Concentrations by UV/F | 64 |
| 3.2.2.1b Oil Composition by GC2 | 64 |
| 3.2.2.1c Aromatic Hydrocarbons Compositions by GC²/MS ... | 64 |
| 3.2.2.2 Benthic Transects | 64 |
| 3.2.2.2a Oil Concentrations by UV/F | 64 |
| 3.2.2.2b Oil Composition by GC2 | 64 |
| 3.2.2.2c Aromatic Hydrocarbon Composition by GC2/MS | 67 |
| 3.2.2.3 Bay 9 Beach . . .0.0...0 . . , ..00.... .0.,,.,,0. | 67 |
| 3.2.3 Bay70 , *...* | 67 |
| 3.2.3.1 Tissue Plots . 0....00.0 | 67 |
| 3.2.3.1a Oil Concentrations by UV/F | 67 |
| 3.2.3.1b Oil Composition by GC² | 70 |
| 3.2.3.1c Aromatic Hydrocarbon Composition by GC²/MS | 70 |
| 3.2.3.2 Benthic Transects | 70 |
| 3.2.3.2a Oil Concentrations by UV/F | 70 |
| 3.2.3.2b Oil Composition by GC²0..... .0..... | 70 |
| 3.2.4 Milne Inlet Samples . ..0...0..0...0.. | 70 |
| 3.2.3.3a Oil Concentrations by UV/F | 73 |
| 3.2.3.3b Oil Composition by GC2 | 73 |
| 3.2.3.3c Aromatic Hydrocarbon Composition by GC2/MS | 73 |
| 3.2.5 Comparison of UV/F-Derived Petroleum Concentrations and GC²-Derived Results*...* ,0.0... .0.. | 73 |
| 3.2.6 Reanalysis of 1982 Field Samples00...00 | 79 |

TABLE OF CONTENTS
(Continued)

| | <u>Page</u> |
|---|-------------|
| 3.3 Oil in Marine Organisms ** . . . 0.0.00 .O . . . * . * .0 | 79 |
| 3.3.1 <u>Mya truncata</u> | 79 |
| 3.3.1.1 Bay 11 * | 79 |
| 3.3.1.1a Oil Concentrations by UV/F | 79 |
| 3.3.1.1b Oil Composition by GC² | 79 |
| 3.3.1.1c Aromatic Hydrocarbon Composition by GC²/MS | 83 |
| 3.3.1.2 Bay 9 | 83 |
| 3.3.1.2a Oil Concentrations by UV/F | 83 |
| 3.3.1.2b Oil Composition by GC² | 85 |
| 3.3.1.2c Aromatic Hydrocarbon Composition by GC²/MS | 85 |
| 3.3.1.3 Bay 7 | 85 |
| 3.3.1.3a Oil Concentrations by UV/F | 85 |
| 3.3.1.3b Oil Composition by GC² | 85 |
| 3.3.1.3c Aromatic Hydrocarbon Composition by GC²/MS | 85 |
| 3.3.2 <u>Serripes groenlandicus</u> | 85 |
| 3.3.2.1 Bay 11 * | 85 |
| 3.3.2.1a Oil Concentrations by UV/F | 85 |
| 3.3.2.1b Oil Composition by GC² | 88 |
| 3.3.2.1c Aromatic Hydrocarbon Composition by GC²/MS | 88 |
| 3.3.2.2 Bay 9 | 88 |
| 3.3.2.2a Oil Concentrations by UV/F | 88 |
| 3.3.2.2b Oil Composition by GC² | 88 |
| 3.3.2.2c Aromatic Hydrocarbon Composition by GC²/MS | 88 |
| 3.3.2.3 Bay 7 | 91 |
| 3.3.2.3a Oil Concentrations by UV/F | 91 |
| 3.3.2.3b Oil Composition by GC² | 91 |
| 3.3.2.3c Aromatic Hydrocarbon Composition by GC²/MS | 91 |
| 3.3.3.1 Bay 11 * | 91 |
| 3.3.3.1a Oil Concentrations by UV/F | 91 |
| 3.3.3.1b Oil Composition by GC² | 91 |
| 3.3.3.1c Aromatic Hydrocarbon Composition by GC²/MS | 91 |

TABLE OF CONTENTS
(Continued)

| | <u>Page</u> |
|---|-------------|
| 3.3.3.2 Bay9 | 96 |
| 3.3.3.2a Oil Concentrations by UV/F | 96 |
| 3.3.3.2b Oil Composition by GC ² | 96 |
| 3.3.3.2c Aromatic Hydrocarbon Composition by GC ² /MS | 96 |
| 3.3.3.3 Bay700.....* | 96 |
| 3.3.3.3a Oil Concentrations by UV/F | 96 |
| 3.3.3.3b Oil Composition by GC ² | 96 |
| 3.3.3.3c Aromatic Hydrocarbon Composition by GC ² /MS | 96 |
| 3.3.4 <u>Astarte borealis</u>,.....,.,..... | 98 |
| 3.3.4.1 Bay 11 | 98 |
| 3.3.4.1a Oil Concentrations by UV/F | 98 |
| 3.3.4.1b Oil Composition byGC ² | 98 |
| 3.3.4.1c Aromatic Hydrocarbon Composition by GC ² /MS | 98 |
| 3.3.4.2 Bay9 000000. | 98 |
| 3.3.4.2a Oil Concentrations by UV/F | 98 |
| 3.3.4.2b Oil Composition byGC ²0..... | 98 |
| 3.3.4.2c Aromatic Hydrocarbon Composition by GC ² /MS | 103 |
| 3.3.4.3 Bay70 | 103 |
| 3.3.4.3a Oil Concentrations by UV/F0..... | 103 |
| 3.3.4.3b Oil Composition byGC ²0..0. | 103 |
| 3.3.4.3c Aromatic Hydrocarbon Composition by GC ² /MS | 103 |
| 3.3.5 <u>Strongylocentrotus droebachiensis</u>0. | 103 |
| 3.3.5.1 Bay 110..... 00 0..... | 103 |
| 3.3.5.1a Oil Concentrations by UV/F0.....0 | 103 |
| 3.3.5.1b Oil Composition byGC ² | 106 |
| 3.3.5.1c Aromatic Hydrocarbon Composition by GC ² /MS | 106 |
| 3.3.5.2 Bay90.....0.....0000 | 106 |
| 3.3.5.2a Oil Concentrations by UV/F0.0* | 106 |
| 3.3.5.2b Oil Composition byGC ²00.....0..... | 106 |
| 3.3.5.2c Aromatic Hydrocarbon Composition by GC ² /MS0..... | 109 |

TABLE OF CONTENTS (Continued)

| | <u>Page</u> |
|--|-------------|
| 3.3.5.3 Bay7 | 109 |
| 3.3.5.3a Oil Concentrations by UV/F | 109 |
| 3.3.5.3b Oil Composition by GC ² | 109 |
| 3.3.5.3c Aromatic Hydrocarbon Composition by GC ² /MS | 109 |
| 3.3.6 Milne Inlet Animals | 109 |
| SECTION FOUR SHORELINE STUDY | 118 |
| SECTION FIVE DISCUSSION RESULTS | 132 |
| SECTION SIX REFERENCES | 138 |

LIST OF FIGURES

| | |
|---|----|
| Figure 2.1. Location of Cape Hatt, Baffin Island0.... . ..*...* | 4 |
| Figure 2.2. Detail of Test Bay Locations | 5 |
| Figure 2.3. General Description of Benthic Sampling Grid Used in Each Bay | 6 |
| Figure 2.4. Schematic of Analytical Strategy | 9 |
| Figure 2.5. BIOS Analytical Protocols | 11 |
| Figure 3.1. GC ² Traces of Saturated Hydrocarbons from Bay 11 Slick/Sheen Samples | 26 |
| Figure 3.2. GC ² Traces of Aromatic Hydrocarbons from Bay 11 Slick/Sheen Samples | 27 |
| Figure 3.3. Bay 11 Slick/Sheen Aromatic Profiles by GC ² /MS | 28 |
| Figure 3.4. GC ² Traces 16 Liter Water Sample, Ragged Channel (W4005): A-Saturates; B-Aromatics; *-Low Molecular Weight Aromatics | 30 |
| Figure 3.5. GC ² Traces of Large Volume Water Sample, Ragged Channel (L4005): A-Saturates; B-Aromatics; *-Low Molecular Weight Aromatics... .. | 33 |

TABLE OF CONTENTS
(Continued)

LIST OF FIGURES
(Continued)

| | | |
|--------------|---|----|
| Figure 3.21. | GC² Trace of Bay 11 Beach Sediments: A-Saturates S4146 (Unweathered, Undegraded); B-Saturates S4139 (Weathered and Degraded); *-Unidentified Cyclic Alkanes (Terpenoids) | 62 |
| Figure 3.22 | Bay 9Sediment Petroleum Hydrocarbon Content; By UV/F (August 15, 1983) * | 63 |
| Figure 3.23. | GC² Trace of Bay 9 Tissue Plot Sediment: A-Saturated Hydrocarbons; B-Aro matic/Unsaturated Hydrocarbons* | 65 |
| Figure 3.24. | GC ² Trace of Representative Bay 9 Beach Sediment; Saturated Hydrocarbons | 67 |
| Figure 3.25. | Bay 7Sediment Petroleum Hydrocarbon Content ; By UV/F (August 14, 1983) | 68 |
| Figure 3.26. | GC2Trace Bay7 Sediments: A-Saturates Tissue Plot No. 4; B-Saturates Benthic Transect | 69 |
| Figure 3.27. | Aromatic Profiles by GC²/MS of Bay7 Sediments | 71 |
| Figure 3.28. | UV/F Spectra of Milne Inlet Sediment Sample Dilution Series, Showing No Oil Present.. . . . | 72 |
| Figure 3.29. | GC² Trace Milne Inlet Sediment Saturated Hydrocarbons; *-Cycloalkenes | 75 |
| Figure 3.30. | Aromatic Profiles by GC²/MS Milne Inlet Sediment Sample | 76 |
| Figure 3.31. | Summary of Oil Concentrations in <u>Mya truncata</u> , by UV/F, ($\mu\text{g/g}$ dry wt.) | 80 |
| Figure 3.32. | <u>Mya truncata</u> Saturated Hydrocarbon GC ² Determinations | 81 |
| Figure 3.33. | <u>Mya</u> Aromatic Profiles by GC²/MS (Bay 11) | 83 |
| Figure 3.34. | Summary of Oil Concentrations in <u>Serripes groenlandicus</u> , UV/F, ($\mu\text{g/g}$ dry wt.) | 85 |
| Figure 3.35. | UV/F Spectra of <u>Serripes</u> Sample Extract From Bay 11 Combined F2 Fraction | 86 |
| Figure 3.36. | <u>Serripes</u> Saturated Hydrocarbon GC² Determinations | 87 |

TABLE OF CONTENTS
(Continued)

LIST OF FIGURES
(Continued)

| | | |
|--------------|---|-----|
| Figure 3.37. | <u>Serripes</u> UV/F Spectra From J3ays9(A) and 7(B) Showing Non-Petroleum Spectral Interference which was eliminated by Column Chromatography | 89 |
| Figure 3.38. | Summary of Oil Concentrations in <u>Macoma calcaria</u> , by UV/F, ($\mu\text{g/g dry wt.}$) | 91 |
| Figure 3.39. | <u>Macoma</u> UV/F Spectra | 92 |
| Figure 3.40. | Representative <u>Macoma</u> GC² Determinations | 93 |
| Figure 3.41. | <u>Macoma</u> Aromatic Profiles by GC ² /MS (Bal I) | 94 |
| Figure 3.42. | <u>Macoma</u> Saturated Hydrocarbons from Bay 7 | 96 |
| Figure 3.43. | UV/F Spectra of <u>Astarte</u> From Bay 11 | 97 |
| Figure 3.44. | Summary of Oil Concentrations in <u>Astarte borealis</u> , by UV/F, ($\mu\text{g/g dry wt.}$) | 98 |
| Figure 3.45. | <u>Astarte</u> Saturated Hydrocarbon GC² Determinations | 100 |
| Figure 3.46. | <u>Astarte</u> Aromatic Profiles by GC²/MS (Bal I) | 101 |
| Figure 3.47. | Bay 11 Urchin Results: A-UV/F; B-Saturated Hydrocarbon GC ² ; B-Aromatic/Unsaturated Hydrocarbon GC ² | 103 |
| Figure 3.48. | Summary of Oil Concentrations in <u>Strongylocentrotus droebachiensis</u> , by UV/F, ($\mu\text{g/g dry wt.}$) | 104 |
| Figure 3.49. | <u>Strongylocentrotus</u> Aromatic Profiles by GC²/MS (Bay 11) | 106 |
| Figure 3.50. | <u>Strongylocentrotus</u> UV/F Spectra , Bay | 107 |
| Figure 3.51. | GC² Trace of <u>Strongylocentrotus</u> ; Bay 7 Saturated Hydrocarbons | 109 |
| Figure 3.52. | Bay 11 Sediment Oil Concentrations (1980-1983) ($\mu\text{g/g}$; by UV/F) | 111 |
| Figure 3.53. | Oil Concentrations in Animals: Bay 11 ($\mu\text{g/g}$; by UV/F) (1980-1983) | 112 |

TABLE OF CONTENTS
(Continued)

LIST OF FIGURES
(Continued)

| | <u>Page</u> |
|--|-------------|
| Figure 3.54. Bay 9 Sediment Oil Concentrations (1980-1983) ($\mu\text{g/g}$; by UV/F) | 113 |
| Figure 3.55. Oil Concentrations in Animals: Bay 9 ($\mu\text{g/g}$; by UV/F) (1980-1983) | 114 |
| Figure 3.56. Oil Concentrations in Benthic Animals: Bay 7 ($\mu\text{g/g}$; by UV/F) (1980-1983) | 116 |
| Figure 3.57. Oil Concentrations in Animals: Bay 10 ($\mu\text{g/g}$; by UV/F) (1980-1982) | 117 |
| Figure 4.1. Representative Shoreline Sediment Saturated Hydrocarbon GC² Determinations: A-S4052, LI; B-4057, TI; C-4309, IDBC | 123 |
| Figure 4.2. Representative Shoreline Sediment Saturated Hydrocarbon GC² Determinations: A-4124, CC; B-4120, CC ..* | 124 |
| Figure 4.3. Representative Shoreline Sediment Saturated Hydrocarbon GC² Determinations: A-S4330; B-S4332; C-S4340 ...* | 125 |
| Figure 5.1. Summary of Comparative Fates of Oil from the BIOS Spills | 136 |

LIST OF TABLES

| | |
|--|----|
| Table 1.1. Hydrocarbon Biogeochemistry (Year 4) Goals | 2 |
| Table 2.1. UV Spectrofluorometry Analytical Conditions | 16 |
| Table 2.2. Fused Silica Capillary Gas Chromatography/Flame Ionization Detection Analytical Conditions | 18 |
| Table 2.3. Explanation of Petroleum Weathering and Source Ratios | 19 |
| Table 2.4. Saturated and Aromatic Hydrocarbon Parameters of Lago Medio Crude Oil | 20 |
| Table 2.5. Gas Chromatography/Mass Spectrometry Instrumental Conditions | 22 |
| Table 2.6. Gas Chromatography/Mass Spectrometry Analytical Outputs | 23 |
| Table 3.1. Seawater Analytical Results (15 Liter) | 25 |
| Table 3.2. Seawater Analytical Results (Large Volume Samples) | 32 |

TABLE OF CONTENTS
(Continued)

LIST OF TABLES
(Continued)

| | <u>Page</u> |
|---|-------------|
| Table 3.3. Summary of Bay 11 Sediment Hydrocarbon Concentration Data | 35 |
| Table 3.4. Bay 11 Sediment Hydrocarbon Compositional Data by GC2 . * ..*... 38 | 38 |
| Table 3.5. Summary of Microbiology Sediment Sample Hydrocarbon Data | 45 |
| Table 3.6 Bay 11 Sediment Core GC2 Data**.. | 51 |
| Table 3.7. Bay 11/12 Deep Water Dredges Sediment Petroleum Hydrocarbons by UV/F . ..0..... *..... ●● | 57 |
| Table 3.8. Shoreline Study 1983 Hydrocarbon Chemistry Results - Analytical Results; Ragged Channel Beaches | 61 |
| Table 3.9. Milne Inlet Subtidal Sediment Control Samples - Petroleum Hydrocarbon Content by UV/F**.. | 73 |
| Table 3.10. UV/F Versus GC ² - Derived Results on Sediments | 77 |
| Table 3.11. QC/QA Analyses: Comparison of Results in 1982 Field Samples . . . | 79 |
| Table 4.1. Shoreline Study Hydrocarbon Chemistry Results Sample Descriptions | 119 |
| Table 4.2. Shoreline Study 1983 Hydrocarbon Chemistry Results - Analytical Results; 1980 Test Plots0.. | 126 |
| Table 4.3. Shoreline Study 1983 Hydrocarbon Chemistry Results - Analytical Results; 1981 Test Plots | 127 |
| Table 4.4. Shoreline Study 1983 Hydrocarbon Chemistry Results - Analytical Results; 1982 Test Plots*...* | 128 |
| Table 4.5. Shoreline Study 1983 Hydrocarbon Chemistry Results - Analytical Results; Ragged Channel Beaches*...*0. | 129 |
| Table 4.6. Shoreline Study 1983 Hydrocarbon Chemistry Results - Analytical Results; Norwegian Test Plots*.. | 130 |



EXECUTIVE SUMMARY

The fourth year of a continuing series of analytical chemical studies of oil fate and transport from the Baffin Island Oil Spill (BIOS) program has been undertaken. Weathered oil continues to erode off the Bay 11 (untreated oil spill test site) beach resulting in increasing oil levels in the Bay 11 sediment. An overall sixfold concentration increase (up to 410 ppm) has been detected in the 3 meter and 7 meter sediments with concentrations decreasing offshore. Transport of oil residues to the deeper areas (35 m) of the Bay 11/12 Area (1-8 ppm) has been detected. Oil in the sediments is more highly weathered than was observed in 1982, although pockets of relatively fresh oil remain on the Bay 11 beach.

Oil concentrations in the sediments of Bay 9 after reaching a high of ~10 ppm in 1981 decreased to 1-3 ppm in 1982, but were seen to increase to levels of 5-10 ppm in 1983.

Detrital feeding benthic animals in Bay 11 appear to have achieved a balance of uptake and depuration of oil while decreasing (through metabolism?) their toxic aromatic hydrocarbon burden. Highest levels of oil are found in Macoma (~60 ppm) and urchins (100 ppm) from Bay 11. Oil was detected in urchins at similar concentrations in Bay 9 although values for the other animals were much lower. Bay 7 remains relatively unimpacted.

Levels of oil in the water column are very low, generally less than 0.5 ppb. Correspondence of the 16 Liter and large volume water samples is fair with levels being somewhat lower (factor of 2) in the large volume samples. Oil at concentrations of ~1 ppb in was detected in Ragged Channel both types of samples with the compositions being similar.

Levels of aromatic hydrocarbons were lower in both the sediments and animals and were not directly related to absolute UV/F-determined oil levels.

Correspondence of UV/F with GC/GCMS data is very weak as was observed in the 1982 study. However, the UV/F signal should be considered to be most useful on a comparative basis as the detection of individual saturated and aromatic components becomes more difficult.

Continuing and increasing oil introduction to Bay 11 sediments caused by erosion of oil off the Bay 11 beach, and transport of sediment offshore are likely future scenarios in light of the above findings.



SECTION ONE

INTRODUCTION

1.1 Project Goals

The analytical chemistry component of the Baffin Island oil spill (BIOS) project involved two major tasks during the fourth year of the project:

1. Nearshore Study - Establishing the concentrations of residual oil, its transport paths, fates, and weathering in the three bays (Bay 11, 9, 7) in the various basic environmental compartments (i.e., water column, benthic sediments, organisms, shoreline) from samples of these compartments taken during the summer of 1983.
2. Shoreline Study - Performing chemical measurements of the oiled shoreline plots to determine concentration and composition of residual oil.

As in previous years (see Boehm et al., 1982a,b; Boehm, 1981; 1983a, b), a tailored analytical program combining analytical property measurements [i.e., ultraviolet fluorescence (UV/F) to determine oil concentrations in the various environmental components] with detailed compositional measurements [i.e., fused silica capillary gas chromatography (GC²) combined with and computer-assisted gas chromatographic mass spectrometry (GC²/MS)] was utilized.

The specific goals of the analytical chemistry program are given in Table 1.1.

1.2 Technical Plan

The analytical plan used in this study was nearly identical to that used previously and involved the following sample types: surface sediments, sediment floe, sediment cores, beached sediments, benthic animal tissues (5 species), water column samples. The types of analyses used: UV/fluorescence, capillary gas chromatography and gas chromatographic mass spectrometry were also used previously. The rationale for each type of analytical procedure is presented in detail in Section Two of this report. The overall plan was to carefully blend analytical techniques of varying sophistication and resolution to best enable the program goals to be achieved within budgetary constraints. Such blends have been successfully employed previously in this and other programs.

TABLE 1.1. HYDROCARBON BIOGEOCHEMISTRY (YEAR 4) GOALS

Nearshore (Ragged Channel)

1. To compare the composition and fates of oil as it impacted the three bays (11, 9, 7).
2. To examine the composition and concentration of high molecular weight petroleum components (n-C10 to n-C32; alkylated benzenes to perylene) in a limited set of water column samples taken from the four bays.
- 3* To examine the chemical nature and weathering of residual surface slick oil and beached oil.
4. To examine the composition and concentration of oil in bottom sediments.
5. To analyze bottom sediments from Bays 11, 9 and 7 for oil content, composition, and weathering changes; to examine the relation of bulk sediment hydrocarbon chemistry; to that of the deposited surface flocculent layer in Bay 11; to examine possible trends in biodegradation; to examine possible offshore transport of the oil.
6. To examine the deputation of petroleum residues by several species of benthic marine organisms, and to examine how these processes varied by species and by bay.
7. To analyze a set of sediment samples from the Milne Inlet subtidal sediments (unimpacted area).

Shoreline Study (Z:Lagoon Area)

1. To determine the concentration and composition of residual oil remaining from several sets of shoreline oil spill countermeasure experiments.
-
-

SECTION TWO

SAMPLING AND ANALYTICAL METHODS

2.1 Sampling

Samples of seawater, offshore **subtidal** sediments, sediment cores, beach sediments, and **benthic** animals were collected from the experimental bays on Cape Hatt, Baffin Island, during August, 1983 (Figures 2.1, 2.2). Bay 11 had been the site of the untreated surface oil spill; Bay 9 had been the site of the chemically dispersed oil spill (Figure 2.2). A detailed description of the sampling techniques used appears in Boehm (1981a) and Boehm et al. (1982a). A brief summary of the **sampling** design and methodology is repeated here.

The sediment and tissue sampling design centered around the grid shown in Figure 2.3 which was identical to that used in 1981 and 1982 collections. Sampling activities occurred during one time during which a large amount of **oversampling** took place (**vis-a-vis** number of samples eventually analyzed).

In August a complete surface sediment collection (tissue plots and **benthic** transects) was obtained from each bay as was a complete collection of the five **benthic** species (Mya truncata, Serripes groenlandicus, Macoma calcarea, Astarte borealis, Strongylocentrotus droebachiensis). Surface floe was obtained from the tissue plots in Bay 11. Sediment cores (0- 15 cm) were obtained at the north and south ends of the 3 and 7 m stratum in Bay 11 only. Several deep sediment samples were also obtained in the Bay 11/ 12 area further offshore in 35 m of water. Sediment samples were obtained at two microbiology transects in Bays 11 and 7 during August.

The water column sampling design is described in Humphrey et al. (1983) and the shoreline sampling design in Owens et al. (1983).

2.1.1 SEAWATER SAMPLING

Two parallel sets of seawater samples were collected. These consisted of 16 liter whole water samples, and large volume samples (~ 100 liters). The water samples




FIGURE 2.1. LOCATION OF CAPE HATT, **BAFFIN** ISLAND.




FIGURE 2.2. DETAIL OF TEST BAY LOCATIONS.




FIGURE 2.3. GENERAL DESCRIPTION OF **BENTHIC** SAMPLING GRID USED IN EACH **BAY**.

were collected **mainly** in Bay 11 although samples were obtained from Ragged Channel Milne Inlet and two from Bay 7 (The reader is referred to Humphrey, 1984 for station locations).

A pumping system was used to collect the seawater samples. A 4-liter solvent-rinsed glass bottle was filled with seawater (four times at each station), sealed with a sheet of Teflon and a screw cap, and stored at ambient temperatures 'until transported to the field laboratory (within 8 hours). At the field laboratory, the samples were preserved by adding 75 ml of Freon 113 to the bottle and then stored **at** room temperature until extraction.

Samples for large-volume high-molecular-weight hydrocarbon analysis were collected with an in situ filtration/absorption sampler. The sampler consisted of a submersible pump, a 293-mm glass fiber filter held in a stainless steel holder, a series of polyurethane plugs in a glass cylinder held in a Teflon sleeve and a flow measurement device. The apparatus was deployed for a period of 4 to 12 hours during which ~100 liters of seawater were pumped through the sampler. Particulate in the seawater were trapped on the filter which was simply folded, placed in an aluminum foil pouch and frozen. Dissolved organics were adsorbed to the polyurethane plugs in the glass cylinder which was sealed on each end with a sheet of Teflon and frozen.

2.1.2 SEDIMENT SAMPLING

Sediments were collected from the beaches in Bay 9, Bay 11 and the countermeasures test area (shoreline study) and from the **subtidal** bottom in Bays 9, 11 and 7 for high-molecular-weight hydrocarbon analysis. Beach sediment stations were located using transect markers established in Bay 9 and Bay 11 and from beach plot markers in the counter-measures test area. The samples from the 1980, 1981, 1982 countermeasures plots (shoreline study in Z-lagoon) were taken from randomly predesignated subareas within a test plot. Beach sediments from Bays 9 and 11 were sampled from a variety of surface and subsurface locations on each beach.

At each station, beach material was scooped into a solvent-rinsed glass jar with a stainless steel trowel. Surface sediment was taken from the top 5 centimeters, subsurface sediment from a depth of 10-15 cm. Care was taken to ensure that the subsurface sample was not contaminated with surface sediment. The samples were transported to the field laboratory and frozen.

Divers collected offshore surface sediment (0-2 cm) by scooping a glass jar along the sediment surface. Unfilled jars were taken through the water surface in a PVC tube whose ends were capped with PVC screw caps and sealed with polyethylene bags. Once below the surface the bags were cut, allowing the tube to flood with seawater and become negatively buoyant. Jars were dispensed from the bottom of the tube and replaced at the top of the tube when filled with sediment.

Divers collected sediment floc with a sampler that consisted of an inverted polyethylene funnel (diameter = 20 cm), a length of Tygon tubing (1 cm diameter x 1 m length), a submersible pump, a metal diverter valve and a stainless steel filter holder (142 mm diameter). The collection procedure is described in **Boehm, 1983a**.

2.1.3 BENTHIC ANIMAL SAMPLING

Benthic animals were collected from Bays 11, 9 and 7.

Divers picked *Mya truncata* and *Strongylocentrotus droebachiensis* using clean gloves. Animals collected from individual stations were placed in nylon mesh bags which were sealed in plastic bags underwater before being carried through the water surface. The contents of the mesh bag were transferred to a plastic bag, labeled, and transported to the field laboratory. The animals were then sorted by species, wrapped in aluminum foil, and frozen.

Other species *Macoma calcaria*, *Astarte borealis* and *Serripes groenlandicus* were airlifted from the sediment by divers. The airlift transferred animals, rocks and mud from the sediment surface into a mesh bag at the opposite end of the airlift. The mesh bag was carried through the water surface in a plastic bag and transported to the field laboratory. The animals were picked from the agglomeration of debris, sorted by species, wrapped in aluminum foil, and frozen.

2.2 Analytical Methods

The general analytical strategy for the chemical assessment consisted of three levels (Figure 2.4). In the first level, samples were extracted and analyzed by ultraviolet spectrofluorometry (UV/F) to measure the concentration of petroleum. Those samples either containing high levels of petroleum or of interest due to sampling time and position




FIGURE 2.4. SCHEMATIC OF ANALYTICAL STRATEGY.

were carried through to the second level, fused silica glass capillary gas chromatography with flame ionization detection (**GC²**). This technique was used to quantify hydrocarbons, to distinguish petroleum hydrocarbons from biogenic hydrocarbons, and to evaluate the composition of petroleum. Measurement of levels of individual aromatic hydrocarbons was **accomplished** during the third phase when computer-assisted **gas-chromatographic/mass spectrometry (GC²/MS)** was used.

The basic types of **samples** (water, sediments, and tissues) were **analyzed** within this study, each according to a slightly different analysis scheme. Each sample type required a unique initial **processing/sample** extraction protocol and followed its own analytical scheme (see Figure 2.5). All samples were spiked with internal standards, androstane (saturated hydrocarbons) and **o-terphenyl** (aromatic hydrocarbons) prior to solvent extraction.

2.2.1 SAMPLE PROCESSING

2.2.1.1 Water Samples (16 Liters)

Sixteen-liter seawater samples were analyzed for high molecular weight hydrocarbons by GC². The water was processed in the field laboratory by extracting three times with Freon. The three extracts were combined, reduced in volume to 10 ml by rotary evaporation and transferred to a glass tube for shipment. Procedural blanks were processed periodically to check for contamination during the field processing.

When received at **Battelle**, the extracts were dried with sodium sulfate, evaporated to <1 ml by rotary evaporation, and displaced with hexane. Three micrograms of two internal standards, androstane and **o-terphenyl**, were added to the extract. An **aliquot** of the extract was weighed on a Cahn Model 25 **electrobalance** to determine total extractable organics. All samples contained large amounts of total extractable organics (natural lipids) and were therefore fractionated by silica gel/alumina column chromatography (see Boehm et al., 1982a) into saturated and unsaturated/aromatic fractions which were analyzed by GC² (see Boehm et al., 1982a). Aromatic fractions and total extracts of selected samples were analyzed by **GC²/MS** (see Boehm et al., 1982a).




FIGURE 2.5. BIOSANALYTICAL PROTOCOLS.

2.2.1.2 Large Volume Water Samples (90-100 Liters)

Each large volume water sample consisted of a glass fiber filter containing particulate **organics** and a polyurethane plug containing dissolved **organics**, both of which were **analyzed** for high-molecular weight hydrocarbons by GC². The filters were processed by cutting them into small pieces which were placed into 250-ml Teflon jars. Three micrograms of two internal standards (androstane and o-terphen yl) and 100 ml of a mixture of dichloromethane and methanol (9:1) were added. The jars were shaken for four hours, and the solvent was decanted. The extraction was repeated with two additional portions of solvent, and the three extracts were combined.

The plugs were processed by squeezing them in the presence of methanol followed by extracting them in a Soxhlet extractor for 24 hours with methanol to remove water and then with **dichloromethane**: methanol (9: 1) to extract organic compounds. All solvent extracts from a sample were combined in a one-liter separator funnel, the **dichloromethane** layer was drawn off, and the remaining water/methanol was extracted three times with 75 ml of **dichloromethane**. The **dichloromethane** extracts from a sample were combined with the combined filter extracts to yield one combined "**dissolved**" plus particulate hydrocarbon sample. The combined extracts reduced in volume to < 1 ml by rotary evaporation and displaced with hexane. An **aliquot** of each of the extracts was weighed on a Cahn Model 25 **electrobalance** to determine total extractable organics. The extracts were fractionated by silica gel/alumina column chromatography (see Section 2.2.2.2) into saturated and unsaturated/aromatic fractions which were analyzed by capillary GC². Aromatic fractions of selected samples were analyzed by GC²/MS.

2.2.1.3 Sediment Sample Processing

Three types of sediment samples were collected and analyzed: surface sediment samples (0-2 cm), oiled beach sediments and, surface floe samples. The sediment and floe were analyzed by different protocols.

2.2.1.3a Surface Sediments (**Benthic** Transects, Tissue Plots, **Microbiology** Sediments, Sediment Cores). Surface sediment samples from the benthic transects and tissue plots were analyzed for high-molecular-weight hydrocarbons using UV/F. One hundred gram **subsamples** were analyzed by UV/F using the analytical method described

below. Selected samples from individual tissue plots and **benthic** transects and all microbiology sediment samples were analyzed by GC².

The extraction method for the sediment UV/F and GC² analysis of sediment samples was based on methods of Brown et al. (1979) and Boehm et al. (1981). Approximately 100 g of wet sediment was weighed into a 250-ml Teflon jar and dried by extracting three times with 75 ml of methanol. Five micrograms of two internal standards, **androstande** and **o-terphenyl** were added to the sediment. The dry sediment was then extracted three times with 100 ml of dichloromethane: methanol (9:1) by shaking on a platform shaker for a minimum of 4 hours for each extraction. All solvent extracts were transferred into a 1-liter separator funnel containing 100 ml of water (Millipore RO) and acidified to a pH of 2 with hydrochloric acid. The dichloromethane layer was drawn off and the aqueous methanol phase was extracted 3 times with 50 ml of dichloromethane. The dichloromethane extracts from a sample were combined, reduced in volume to 1 ml by rotary evaporation and displaced with methanol. The extract was transferred to a 50 ml glass tube containing 10 ml of methanol and 4 ml of 10N aqueous KOH, sealed with a Teflon cap and heated at 800C for 4 hours to saponify interfering polar compounds. The mixture was cooled then extracted 3 times with 15 ml of hexane. The combined hexane extracts were dried over sodium sulfate and concentrated by rotary evaporation to approximately 1 ml. The extract was then analyzed by UV/F (see Section 2.2.2.1).

Single aliquots of extracts to be further analyzed by GC² and/or GC²/MS were weighed on a Cahn Model 25 **electrobalance** to determine total extractable organics. The extracts were fractionated by silica gel/alumina column chromatography into saturated and unsaturated/aromatic fractions which were analyzed by GC². Aromatic fractions of selected samples were analyzed by capillary GC²/MS.

2.2.1.3b Surface Floe Analysis. Surface floe samples were analyzed for high-molecular-weight hydrocarbons using UV/F. A selected subset were analyzed by GC² and GC²/MS techniques. The glass fiber filters containing the floe were extracted with dichloromethane: methanol (9: 1) using the techniques described for the large volume water sample filters. The total extracts were freed of polar compounds which interface with the UV/F by saponification as described for surface sediments. All sample extracts were analyzed by UV/F and selected samples were fractionated by silica gel/alumina column chromatography into saturated and unsaturated aromatic fractions which were analyzed by GC². Selected aromatic fractions were analyzed by GC²/MS.

2.2.1.3c Oiled Beach Sediment Analysis. Oiled beach sediments were analyzed for high molecular weight hydrocarbons using only GC2 techniques. The analytical methodology was the same as that described for GC2 analysis of surface sediments.

2.2.1.4 Benthic Animal Tissue Processing

Five species of benthic bivalves were analyzed: Mya truncata, Serripes groenlandicus, Macoma calcaria, Astarte borealis, and Strongylocentrotus droebachiensis (sea urchin). Samples from individual tissue plot stations were analyzed by UV/F. Subsequently, extracts from all five tissue plot stations at a given depth and bay were combined and analyzed by GC².

The extraction and analytical procedure (see Boehm et al., 1982a) was based closely on that of Warner (1976) as revised by Boehm et al. (1982c). Clam tissues (guts, muscle, gills) were removed from the shells with solvent-rinsed utensils. Samples with more than 10 grams wet weight tissue were homogenized with a Polytron tissue homogenizer, and a 10-30 g aliquot was taken for analysis. Otherwise, the entire sample was homogenized. A small aliquot of the tissue homogenate was taken for wet weight/dry weight determination. Tissue was digested overnight with a 5 N aqueous potassium hydroxide and were then extracted in a separator funnel with hexane. Hexane extracts were combined, dried with sodium sulfate and concentrated to 0.5 ml by rotary evaporation. Polar and biogenic compounds which interfered with the UV/F analysis were removed from the extract by alumina column chromatography. One of two sizes of columns, one containing 6.5 g and the other containing 25 g of 7.5% water deactivated alumina, were used depending on the amount of tissue. The column was eluted with 25 ml or 75 ml of hexane/dichloromethane (9: 1) to isolate the saturated> unsaturated and aromatic compounds. The fraction was concentrated and transferred into hexane for UV/F analysis.

After UV/F analysis, the extracts from the tissue plot stations along each depth stratum were combined, concentrated by rotary evaporation and displaced with hexane. The pooled extracts 5 species x 3 bays were fractionated by silica gel/alumina column chromatography into saturated and unsaturated/aromatic fractions which were analyzed by GC2. Aromatic fractions from these fifteen combined samples were analyzed by GC²/MS.

2.2.2 SAMPLE ANALYSIS

2.2.2.1 UV/F Analysis

The synchronous excitation/emission technique used has been extensively described previously (Boehm et al., 1982a). The analytical conditions are shown in Table 2.1. This technique measures aromatic hydrocarbons with a two- to five-ring aromatic structure (Lloyd, 1971). The extract was repeatedly diluted by 50% and reanalyzed until a comparison of two consecutive dilutions indicated that the analysis was done within the linear range of fluorescence response. The intensity of the fluorescence spectra was measured at 350-360 nm which corresponded to a peak maximum present in a Lagomedio Bay 11 reference oil sample. The fluorescence spectra were converted to relative concentration units by comparing the peak height at each wavelength to that of a Bay 11 oil standard curve.

2.2.2.2 Fractionation

Those sediment, tissue, and water samples chosen for GC2 analyses were fractionated by silica gel/alumina column chromatography prior to GC2 analysis. Column chromatography isolated the saturated and aromatic hydrocarbons from the total extract, thereby facilitating the identification and quantification of individual hydrocarbon compounds which were present in the sample extract.

The total extract was charged to a 100% activated silica gel/5 percent deactivated alumina/activated copper (11 g, 1 g, 2 g) chromatography column that was wet-packed in dichloromethane and prepared by eluting with 30 ml each of dichloromethane and hexane. The column was eluted with 18 ml of hexane followed by 21 ml of hexane:dichloromethane (1: 1) to isolate the saturated (f_1) and aromatic/unsaturated (f_2) hydrocarbons, respectively. After concentrating each fraction by rotary evaporation, the total gravimetric concentration was determined by weighing a measured aliquot on a Cahn Model 25 electrobalance.

TABLE 2.1. UVSPECTROFLUOROMETRY ANALYTICAL CONDITIONS

| | |
|--------------------|-------------------------------------|
| Instrument: | Farrand System 3 spectrofluorometer |
| Features: | Corrected excitation |
| Slits: | |
| Excitation: | 2.5 nm |
| Emission: | 5.0 nm |
| Scan Speed: | 50 nm/min |
| Cell: | 10 nm quartz |
| Monochrometers: | <u>Synchronous</u> |
| Excitation: | 225-475 nm |
| Emission: | 250-500 nm |
| Daily Calibration: | Bay 11 Lagomedio oil |
| Quantification: | External calibration curves |

2.2.2.3 GC2 Analysis

GC2 analysis served to identify and quantify the petroleum hydrocarbon compounds present in the sample. The relative concentrations of individual compounds identified the composition of oil present, and the absolute concentrations served as a measure of the amount of oil present. The concentrations of certain compounds were also used to calculate indicator ratios that reveal the type of hydrocarbons present, i.e., biogenic or petroleum.

Each fraction was analyzed by fused silica capillary gas chromatography on a Hewlett Packard 5840 or 5880 gas chromatography equipped with a **splitless** injection port and a flame ionization detector. **Wall** coated open tubular (**WCOT**) fused silica columns (0.25 mm x 30m, J&W Scientific) coated with SE30 and **SE54** stationary phases were used to analyze, respectively, the **f₁** and **f₂** fractions from the column chromatography. The instrumental conditions are listed in Table 2.2. Compounds were identified by comparing retention indices of peaks in the samples to retention indices of known compounds in a standard mixture that was analyzed daily. Concentrations were calculated by comparing the integrated areas of peaks with the area of the appropriate internal standard (androstane for the **f₁**, **o**-terphenyl for the **f₂**). The total concentrations of saturated and aromatic hydrocarbons were determined by planimetering the unresolved area, converting it to integrator area units, adding it to the total resolved integrated area, and calculating a concentration using the internal standard method.

The concentrations of **n-alkanes** and isoprenoids were determined from GC² on a dry weight basis. From these concentrations, a series of key diagnostic parameters were calculated (Tables 2.3 and 2.4). These ratios are useful in establishing the composition of the oil, the contribution of biogenic hydrocarbons, and the degree that the oil was weathered when compared to values for the spilled oil itself (Table 2.4).

Concentrations of the chromatographically unresolved complex mixture (**UCM**) appearing as a “hump” on the GC2 traces were quantified relative to the internal standard by **plainimetry**. This UCM is characteristic of residual petroleum hydrocarbons in the samples.

TABLE 2020 FUSED SILICA CAPILLARY GAS CHROMATOGRAPHY/FLAME
IONIZATION DETECTION ANALYTICAL CONDITIONS

| | |
|-----------------------|--|
| Instrument: | Hewlett Packard 5840 or 5880 gas chromatography |
| Features: | Split/splitless capillary inlet system Microprocessor-controlled functions |
| Inlet: | Splitless |
| Detector: | Flame ionization |
| Column: | |
| f₁: | 0.25 mm I.D. x 30 m SE30 fused silica (J&W Scientific) |
| f₂: | 0.25 mm I.D. x 30 m SE54 fused silica (J&W Scientific) |
| Gases: | |
| Carrier: | Helium 2 ml/min |
| Make-up: | Helium 30 ml/min |
| Detector: | Air 300 ml/min (500 ml/min for 5880) |
| Temperatures: | |
| Injection port: | 250°C |
| Detector: | 300°C |
| Column oven: | 40°-290° 3° C/min |
| Daily calibration: | Alkane/aromatic mixture |
| Quantification: | Internal standards (F₁ androstane; f2 o-terphenyl) |

TABLE 2.3. EXPLANATION OF PETROLEUM WEATHERING AND SOURCE RATIOS

1. The Biodegradation Ratio (Alkane/Isoprenoid)

$$\text{ALK/ISO}_{14-18} = \frac{1400 + 1500 + 1600 + 1700 + 1800}{1380 + 1470 + 1650 + 1708 + 1810}$$

The ALK/ISO ratio approaches 0 as the n-alkanes are depleted.

2. The n-C₁₈/Phytane Ratio

The C₁₈/Phy ratio also approaches 0 as n-C₁₈ is preferentially depleted

3. The Pristane/Phytane Ratio

The Pris/Phy ratio is equal to ~0.7 in aged Lagomedio oil. As the amounts of the biogenic isoprenoid, pristane increase relative to the petrogenic isoprenoid, phytane, this ratio becomes large.

4. The Saturated Hydrocarbon Weathering Ratio (SHWR)

$$\text{SWHR} = \frac{\text{sum of n-alkanes from n-C}_{10} \text{ to n-C}_{25}}{\text{sum of n-alkanes from n-C}_{17} \text{ to n-C}_{25}}$$

The SWHR approaches 1.0 as low-boiling saturated hydrocarbons (n-C₁₀ to n-C₁₇) are lost by evaporation.

5. The Aromatic Weathering Ratio (AWR)

$$\text{AWR} = \frac{\text{Aaryl benzenes} + \text{naphthalenes} + \text{fluorenes} + \text{phenanthrenes} + \text{dibenzothiophenes}}{\text{Total phenanthrenes} + \text{dibenzothiophenes}}$$

The AWR approaches 1.0 as low-boiling aromatics are lost by evaporation and/or dissolution.

6. Carbon Preference Index (CPI)

$$\text{CPI} = \frac{2(\text{n-C}_{27} + \text{n-C}_{29})}{\text{n-C}_{26} + 2\text{n-C}_{28} + \text{n-C}_{30}}$$

CPI \cong 1.0 for petroleum

CPI ranges from 3-6 for terrigenous plant waxes.

The relative amounts of petroleum alkanes to terrigenous biogenics can be assessed through this ratio.

TABLE 2.4. SATURATED AND AROMATIC HYDROCARBON
PARAMETERS OF LAGOMEDIO CRUDE OIL^a

| | Fresh Oil | Aged Oil |
|-------------------------------------|-----------|----------|
| <u>Saturates</u> | | |
| SHWR | 2.9 | 2.3 |
| ALK/ISO | 2.4 | 2.5 |
| Pris/Phy | 0.85 | 0.74 |
| n-C ₁₈ /Phy ^b | 1.6 | 1.6 |
| <u>Aromatics</u> | | |
| AWR | 4.3 | 3.5 |

aKey

$$\text{SHWR} = \frac{(\sim \text{n-alkanes; C}_{10}\text{-C}_{25})}{(\sum \text{n-alkanes; C}_{17}\text{-C}_{25})}$$

$$\text{AWR} = \frac{(\text{Alkyl Benzenes} + \text{Naphthalenes} + \text{Fluorenes} + \text{Phenanthrenes} + \text{Dibenzothiophenes})}{\text{Phenanthrenes} + \text{Dibenzothiophenes}}$$

$$\text{ALK/ISO} = \frac{(\sum \text{alkanes; C}_{14}\text{-C}_{18})}{(\sum \text{5 isoprenoids; in n-C}_{13} \text{ boiling range})}$$

PRIS = pristane

PHY = phytane

b Note: This ratio was expressed as the inverse in previous BIOS reports.
It is reformulated here to be consistent in concept to the ALK/ISO ratio.

2.2.2.4 Gas Chromatography/Mass Spectrometry (GC²/MS)

Selected samples suspected to contain petroleum by the GC2 analyses were analyzed by GC²/MS to measure the concentration and composition of individual aromatic hydrocarbons in the samples. The concentrations of a series of **polynuclear** aromatic hydrocarbons, in particular the **alkylated** phenanthrenes and dibenzothiophenes, serve as a fingerprint of weathered petroleum.

The f2 (aromatic fraction) from the silica gel/alumina column chromatography was analyzed for **polynuclear** aromatic hydrocarbons by GC²/MS. An aliquot of the fraction was analyzed using a Finnegan 4530 instrument equipped with a 0.25 mm x 30 m SE54 fused silica capillary column (J&W Scientific), which was threaded directly into the ion source. Instrumental conditions are listed in Table 2.5.

Selected ion searches were used to obtain ion chromatograms for aromatic compounds with known retention indices and suspected to be present in the samples. Concentrations of the identified compounds were determined by measuring peak areas of the appropriate peaks in the selected ion chromatograms and relating them to that of the internal standard. Relative response factors for each component were calculated from analyses of analytical standards, if available, or were extrapolated. The compounds reported from the GC²/MS analyses are listed in Table 2.6 and are presented in a series of Figures in the results section with compound designations as in Table 2.6.

TABLE 2.5 GAS CHROMATOGRAPHY/MASS SPECTROMETRY
INSTRUMENTAL CONDITIONS

| | |
|--------------------------------|---|
| INSTRUMENT: | Finnegan 4530 gas chromatograph/mass spectrometer |
| FEATURES: | Data General Nova 3 data system with Inc os data system Finnegan MAT 9610 |
| INLET: | Splitless |
| DETECTOR: | Quadruple mass spectrometer |
| SCAN RATE: | 450 amu/sec (45-450 amu) |
| IONIZATION VOLTAGE: | 70 eV |
| COLUMN | 0.25 mm id. x 30 m SE54 fused silica (J&W Scientific) |
| INTERFACE: | Direct insertion of column into source |
| CARRIER GAS: | Helium 2 ml/min |
| TEMPERATURES | |
| INJECTION PORT: | 270°C |
| SEPARATOR OVEN: | 280°C |
| SOURCE: | 250°C |
| GC OVEN: | 40-290°C, 10°C/min (temperature program) |
| DAILY CALIBRATION: | FC43, DFTPP and aromatic mixture |
| QUANTIFICATION: | Internal standard (o-terphenyl) (response factors) |

TABLE 2.6 GAS CHROMATOGRAPHY/MASS **SPECTROMETRY** ANALYTICAL OUTPUTS

| POLYNUCLEAR AROMATIC HYDROCARBONS | |
|--|---|
| Abbreviation | Compound or Compound Grouping |
| AB | Alkyl benzenes (C3 to C₆) |
| N | Naphthalene homologous series |
| C₀N | Naphthalene |
| C₁N | Methyl naphthalenes |
| C₂N-C₄N | Alkyl naphthalenes |
| F | Fluorene homologous series |
| C₀F | Fluorene |
| C₁-C₃F | Alkylated fluorenes |
| P | Phenanthrene homologous series |
| COP | Phenanthrene |
| C₁PC₄P | Alkylated phenanthrenes |
| DBT | Dibenzothiophene homologous series |
| C₀DBT | Dibenzothiophene |
| C₁DBT-C₃DBT | Alkylated dibenzothiophenes |
| | Fluoranthene |
| | Pyrene |
| | Benzo(a)anthracene |
| PAH | Chrysene |
| | Benzo(a)fluoranthene |
| | Benzo(a)pyrene |
| | Benzo(e)pyrene |
| | Perylene |

SECTION THREE

RESULTS (NEARSHORE STUDY)**3.1 Water Column****3.1.1 OIL ON THE WATER'S SURFACE (SURFACE SHEEN SLICK)**

Two 16 Liter samples of seawater were collected in an area of visible sheening in Bay 11. These sheens, which originated from the stranded oil residues on the Bay 11 beach, were analyzed mainly to determine the composition of the surface sheens, their weathering characteristics as well as the total oil concentrations in the 16 Liter surface water sample.

Concentrations of petroleum in these samples (Table 3.1) were 1870 and 1030 $\mu\text{g}/\text{sample}$. These samples represent the source petroleum leached from the Bay 11 beach. The compositional characteristics of the two samples are similar (Figures 3.1 and 3.2). The petroleum is highly weathered having lost much of its n-alkane material in the n-C₁₀ to n-C₁₇ range. Normal alkanes remain prominent in the n-C₂₀ to n-C₃₄ range. SHWR ratios for both samples are -1.0. That the residual oil has been extensively biodegraded is evident from the low C₁₈/Phy ratios (-0.26). The higher molecular weight normal alkanes appear to be exempted from this microbial attack. Similar weathered residues from temperate spills (e.g., Amoco Cadiz, Boehm et al., 1981) exhibit near-total depletion of all n-alkanes throughout the boiling range (see Figure 1.4 in Boehm et al., 1982a).

Surface sheen samples taken in 1982 (see Boehm, 1983a) were weathered to a lesser extent (SHWR = 2.0; C₁₈/Phy = 1.6) than the samples taken in 1983.

Both of these samples were analyzed by GC²/MS. The results are presented in Figure 3.3. The aromatic compositions are comprised entirely of alkylated phenanthrene and dibenzothiophene (DBT) compounds. Sample W 4022 also contained tri and tetramethyl naphthalenes. These compositions convert to the AWR ratios presented in Table 3.1

TABLE 3.1. SEAWATER ANALYTICAL RESULTS (16 LITER)

| Sample ID | Volume (L) | Description | Hydrocarbon Concentrations (µg/L) | | | SHWR C | lg/Phy | AWR | Oil? |
|--------------|------------|---|-----------------------------------|--------------|--------------|--------|--------|------------|-------|
| | | | Saturates | Aromatics | Total | | | | |
| W4021 | NA | Bay 11; Surface Slick 8-12-83; 1300 | 330a | 1540a | 1870a | 1.1 | .27 | 1.0 | Yes |
| W4022 | NA | Bay 11; Surface Slick | 460a | 570a | 1030a | 1.0 | .26 | 1.1 | Yes |
| W4003 | 15.9 | Bay 11; S. Micro; 10m 8-12-83; 1017 | .26 | .11 | .37 | -- | -- | ND | No |
| W4004 | 15.4 | Bay 11; S. Micro; 1m 8-12-83; 1243 | .20 | .33 | .53 | -- | -- | ND | Trace |
| W4008 | 16.1 | Bay 11; S. Micro; 10m 8-13-83; 1540 | <.1 | <.1 | <.1 | -- | -- | -- | No |
| W4009 | 12.2 | Bay 11; S. Micro; 1m 8-13-83; 1949 | <.1 | <.1 | <.1 | -- | -- | -- | No |
| W4010 | 15.8 | Bay 11 ; S. Micro; 10m 8-15-83; 1131 | .14 | .17 | .31 | -- | -- | -- | No |
| W4011 | 16.0 | Bay 11; S. Micro; 1m 8-15-83; 0750 | .18 | .18 | .36 | -- | -- | -- | No |
| W4013 | 15.8 | Bay 11; S. Micro; 10m 8-16-83; 0810 | 1.1 | <.1 | 1.1 | -- | -- | -- | Trace |
| W40 4 | 15.7 | Bay 11; S. Micro; 1m 8-16-83; 1131 | .18 | .10 | .28 | -- | -- | -- | No |
| W40 5 | 15.6 | Bay 11; S. Micro; 10m 8-19-83 ; 0810 | .26 | .1 | .36 | -- | -- | -- | Trace |
| W4016 | 15.2 | Bay 11; S. Micro; 1m 8-19-83; 1207 | .27 | <.1 | .27 | -- | -- | -- | No |
| W 4006 | 15.5 | Bay 7; N. Micro; 5m 8-12-83; 1555 | .26 | .20 | .46 | -- | -- | ND | No |
| W4017 | 15.5 | Bay 7; N. Micro; 5m 8-18-83; 0820 | .17 | <.1 | .17 | -- | -- | -- | No |
| W4005 | 15.8 | Ragged Charnel; 5m 8-16-83; 1735 | 1.7 | 1.0 | 2.7 | -- | -- | -- | Yes |
| W4012 | 16.1 | Milne Inlet; 5m 8-1 5-83; 1521 | <.1 | <.1 | <.1 | -- | -- | -- | No |

a = Values in µg/sample; no volume reported.
 ND= No aromatics detected.




FIGURE 3.1. GC2 TRACES OF SATURATED HYDROCARBONS FROM BAY 11 SLICK/SHEEN SAMPLES.




FIGURE 3.2. GC² TRACES OF AROMATIC HYDROCARBONS FROM BAY 11 SLICK/SHEEN SAMPLES.




FIGURE 3.3. BAY 11 SLICK/SHEEN AROMATIC PROFILES BY GC2/MS.

3.1.2 OIL IN THE WATER COLUMN

3.1.2.1 16 Liter Samples

Analytical results obtained from the 16 Liter water samples are also presented in Table 3.1. Traces of petroleum material diagnosed from the presence of a **petrogenic** series of **n-alkanes** (**n-C₂₀** to **n-C₃₀**) were detected in the saturated hydrocarbon fractions of several of the samples (Figure 3.4). The biogenic **isoprenoid**, pristane, was prominent in nearly **all** of the water samples, much more so than indicated in Figure 3.4A. GC2 analysis of most of the 16 Liter samples failed to indicate the presence of any petrogenic aromatic compounds. Levels of "aromatics" shown in Table 3.1 for the most part are actually indicative of **biogenic** unsaturated hydrocarbon components **eluting** in the **"aromatic" fraction**.

The four samples in which trace quantities ($< 1.0 \mu\text{g/L}$) or greater were detected, include three from Bay 11 (two at 10 meters depth, one at 1 meter depth) and one from Ragged Channel. Levels less than $0.5 \mu\text{g/L}$ represent biogenic material only. The maximum quantity of oil detected, $2.7 \mu\text{g/L}$, was seen in one of the Ragged Channel samples. The composition of this sample is illustrated in Figure 3.4. A small amount of unresolved naphthenic material is observed in the saturated hydrocarbon fraction as well as a prominent series of **n-alkanes** from **C₂₀** to **C₃₃**. The aromatic fraction (Figure 3.4B) contains a series of low boiling components as well as a significant amount of unresolved material as well.

Absolute maximum concentrations in these water samples are similar to those found in a small set of 16 L samples taken in 1982, with **n-alkane** values $\sim 1.0 \mu\text{g/L}$ in both years.

GC²/MS analyses were performed on four additional 16L samples. **No** detectable aromatics (i.e. $> 1.0 \mu\text{g/L}$) were observed in three of the samples. Sample W4005 contained very low levels ($< 1\text{-}5 \text{mg/L}$) of alkylated phenanthrene and DBT compounds.




FIGURE 3.4. GC² TRACES 16 LITER WATER SAMPLE, RAGGED CHANNEL (W4005):
A- SATURATES; B- AROMATICS; *-LOW MOLECULAR WEIGHT AROMATICS.

3.1.2.2 Large Volume Water Samples

The combined dissolved and particulate fractions were analyzed as one water column sample. The results of these analyses are presented in Table 3.2. Although large quantities of biogenic lipids were captured in the samples, cleanup and analyses of the hydrocarbon fractions revealed little evidence of oil in the water column. These samples were obtained in a pair with a 16 L sample in order to compare the quantitative and qualitative results from both sample types. There is generally reasonable agreement between the two sets. The large volume samples contained trace quantities of oil in four samples at levels of 0.1-0.3 $\mu\text{g/Liter}$. Two of the four "trace level" large volume samples from Bay 11 corresponded to paired 16 L samples in which traces of oil were also found (L4004;W4004:L4015; W4015) although the large volume sample levels were about a factor of two to three lower in concentration. The results on the 16 Liter Ragged Channel sample (W4005, Table 3.1), exhibiting higher levels of detected petroleum, were confirmed by the parallel large volume sample (L4005, Table 3.2) although again the large volume sample was lower in concentration by a factor of two. Compositionally, this Ragged Channel large volume sample (Figure 3.5) agreed closely with the 16 L sample results (Figure 3.4). The lower molecular weight components were present in the aromatic fractions from both the large volume sample and the 16L sample.

The 1983 results differed from the previous years results in that the 1983 petroleum residues appeared to be more substantially weathered than those detected in the water column during the 1982 field sampling season. Normal alkanes in the n-C₁₂ to n-C₁₇ boiling range which were frequently detected in the 1982 large volume samples were absent in the 1983 samples. Low levels of naphthalenes detected in the 1982 samples were not seen in the GC2 traces from 1983.


Two samples were analyzed by GC²/MS. No detectable (>1.0ng/L) aromatics were seen in the L4004 or the L4005 sample. The Ragged Channel L4005 sample contained the same low boiling compounds as did the W4005 samples. However, data on these components were not acquired in the GC²/MS analysis, and therefore the nature of these compounds remains unknown.

TABLE 3.2. SEAWATER ANALYTICAL **RESULTS (LARGE VOLUME SAMPLES)**

| Sample ID | Volume (L) | Description | Hydrocarbon Concentrations (µg/L) | | | SHWR | C ₁₈ /Phy | AWR | Oil? |
|-----------|------------|--|-----------------------------------|-----------|-------|------|----------------------|-----|-------|
| | | | Saturates | Aromatics | Total | | | | |
| L4003 | 99 | Bay 11; S. Micro; 10m 8-12-83; 1017 | .09 | .11 | .20 | NA | NA | -- | Trace |
| L4004 | 86 | Bay 11; S. Micro; 1m 8-12-83; 1243 | .23 | .02 | .2s | NA | NA | ND | Trace |
| L4008 | 100 | Bay 11; S. Micro; 10m 8-13-83; 1540 | <.01 | <.01 | <.01 | NA | NA | -- | No |
| W4009 | 109 | Bay 11; S. Micro; 1m 8-13-83; 1949 | <.01 | <.01 | <.01 | NA | NA | -- | No |
| W4010 | 91 | Bay 11; S. Micro; 10m 8-15-83; 1131 | .05 | .07 | .12 | NA | NA | -- | No |
| L4011 | 117 | Bay 11; S. Micro; 1m 8-15-83; 0750 | .02 | .07 | .09 | NA | NA | -- | No |
| L4013 | 90 | Bay 11; S. Micro; 10m 8-16-83; 0810 | .04 | .04 | .08 | NA | NA | -- | NO |
| L4014 | 91 | Bay 11; S. Micro; 1m 8-16-83; 1131 | .07 | .03 | .10 | NA | NA | -- | Trace |
| L4015 | 99 | Bay 11; S. Micro; 10m 8-19-83; 0810 | .10 | .03 | .13 | NA | NA | -- | Trace |
| L4016 | 92 | Bay 11; S. Micro; 1m 8-19-83; 1207 | .08 | .04 | .12 | NA | NA | -- | No |
| L4006 | 90 | Bay 7; N. Micro; 5m 8-12-83; 1555 | .08 | .03 | .11 | NA | NA | -- | No |
| L4017 | 92 | Bay 7; N. Micro; 5m 8-18-83; 0820 | .06 | .03 | .09 | NA | NA | -- | Trace |
| L4005 | 85 | Ragged Channel; 5m 8-16-83; 0735 | .67 | .47 | 1.1 | NA | 0.9 | ND | Yes |
| L4012 | 100 | Milne Inlet; 5m 8-15-83; 1'521 | <.01 | <.01 | <.01 | NA | NA | -- | No |

A= Compounds not detected.

D= No aromatics detected.



3.2 Oil in the Sediments

3.2.1 BAY 11

Six types of sediment samples were analyzed from the Bay 11 beach and **subtidal** region during 1983. Sediment samples (0-2 cm) were **analyzed** from the tissue plots at 3 m and 7 m depth adjacent to the area from which animal samples were acquired for tissue analysis; from the three benthic transects at the same depths in which **benthic** communities were quantified; from the tissue plot surface floe to determine levels and composition of oil in newly deposited sediment; from a microbiologyy transect to examine offshore distributions of oil along a intertidal to 10 meter water depth transect; from a series of deep water sediments to determine transport of oil to deeper areas (~35m) in the Bay 11/12 area; from the Bay 11 beach to determine quantities and composition of stranded oil.

3.2.1.1 Tissue Plots

3.2.1.1a Oil Concentrations by UV/F. Figure 3.6 and Table 3.3 presents a summary of the petroleum concentrations in UV/F-determined Lagomedio oil equivalents in Bay 11 **subtidal** sediments. The Bay 11 sediments contained high levels of **oil**. Concentrations on a log transformed basis [i.e. \bar{X}_G (Lower 95% confidence limit, upper 95% confidence limit)] in the 3 m tissue plots were 22.5 (8.9, 57.8) $\mu\text{g/g}$ compared with 3.0 (1.1, 8.1) $\mu\text{g/g}$ last observed in 1982. The highest value found in 1983, 123 $\mu\text{g/g}$, is only twice the highest 1982 value, 66 $\mu\text{g/g}$. However, on the average **roughly** two times more oil has impacted the three meter stratum in Bay 11. Concentrations at the 7 m stratum [2.6 (0.1, 17.5)] $\mu\text{g/g}$ were lower than at 3 m and were more tightly grouped in concentration. This illustrates the patchy nature of the 3 m distributions caused by active oil deposition at this depth. Further offshore (7 m) distributions are more homogeneous although a factor of three separates the lowest (8.1 $\mu\text{g/g}$) and highest (21.2 $\mu\text{g/g}$) concentration at the 7 m depth. The 1982 sampling revealed levels of oil at 7 m to be 5.3 (2.7, 10.1) $\mu\text{g/g}$, or on the average, a **factor of two lower** than in 1983.

Active erosion of beached oil is occurring in Bay 11 as it was in 1982. Concentrations of oil are higher at the southern end of the 3 m sampling line, as was also the case in 1982.

TABLE 3.3 SUMMARY OF **BAY 11** SEDIMENT HYDROCARBON **CONCENTRATION** DATA

| Sample Type | Depth | Concentration: X (-95%, +95%) ($\mu\text{g/g}$) | |
|---------------------------|------------|--|------------------|
| | | <u>1982</u> | <u>1983</u> |
| Tissue Plot | 3m | 3.0(1.1, 8.1) | 22.5(8.9, 57.8) |
| Tissue Plot | 7m | 5.3(2.7, 10.1) | 12.6(9.1, 17.5) |
| Benthic Transect 1 | 3m | 4.0(2.1, 7.6) | 21.2(8.6, 51.9) |
| | 7m | 6.6(5.4, 8.0) | 22.5(18.7, 27.1) |
| Benthic Transect 2 | 3m | 1.4(.66, 3.1) | 14.0(4.7, 41.3) |
| | 7m | 4.7(3.3, 6.6) | 12.0(10.9, 13.2) |
| Benthic Transect 3 | 3m | 10.3(3.0, 35.0) | 119 (86.7, 162) |
| | 7m | 4.0(2.1, 7.5) | 18.8(9.9, 35.8) |
| Deep Sediment | 35m | ----- | 4.6(2.5, 8.3) |




FIGURE 3.6. BAY 11 SEDIMENT PETROLEUM HYDROCARBON CONTENT; UV/F (AUGUST 13, 1983).

3.2.1.1b Oil Composition by GC². Four samples of Bay 11 tissue plot sediments were analyzed by GC² to determine the oil's composition. These results are summarized in Table 3.4. Two representative saturated hydrocarbon GC² traces are presented in Figure 3.7. A combination of weathered oil (SHWR = 1.1) illustrating a moderate degree of biodegradation (C₁₈/Phy .43) and terrigenous alkanes is illustrated by the GC² trace in Figure 3.4a (tissue plot 8 at 3 meters depth). The UCM material also characterizes this weathered oil residue. Higher concentrations of oil are found (Table 3.4) at tissue plot #10 at 3 meters and the GC² trace of this sample illustrates the same features of weathered oil. The SHWR for this sample is slightly higher (SHWR = 1.3) which indicates that n-alkanes in the C₁₀ to C₁₇ range are still present in this sample. Notice how the increase amount of oil depresses the CPI further (CPI = 1.4) as the relative influence of the terrigenous n-alkanes is less due to the increased amount of oil.

The biodegradation indicator used here, C₁₈/Phy, varies from values of 0.4 to 1.0, indicative of moderately degraded oil, to 1.5 to 2.0 indicative of an undegraded oil.

3.2.1.1c Aromatic Hydrocarbon Composition by GC²/MS. Three tissue plot samples were analyzed by GC²/MS, the results summarized in Figure 3.8. The aromatic hydrocarbon concentration ranges agree well with those determined from the 1982 sample set. Tissue plot 10, however, contained lower concentrations of those quantified homologous series, by a factor of 3, than detected in 1982. The alkylated phenanthrene and DBT series are the most prominent quantifiable aromatic compounds.

3.2.1.2 Surface Floc Samples

3.2. 1.2a Oil Concentrations by UV/F. UV/F determinations on floc samples from each of the Bay 11 tissue plots are presented in Figure 3.6. Concentrations are reported as mg oil/m² assuming a 0.1 m² sampling area as has been used previously. Petroleum levels of .23 to 4.0 mg/m² were observed in the samples. Background levels are ~0.05 mg/m². The 3 m samples had substantially more oil associated with the floc, .93 (.65, 1.3) mg/m², than did the 7 m samples, .29 (.22, .37) mg/m², again indicating the more substantial ongoing impact along the 3m stratum.

The 1981 results had previously indicated that maximum floc impact in the dispersed oil spill (Bay 9) had achieved at 33 mg/m² impact. Bay 11 floc samples taken in 1982 had indicated a lesser but very significant oil impact to the surface floc at 3 m, .54

TABLE 3.4 BAY 11 SEDIMENT HYDROCARBON COMPOSITIONAL DATA BY GC2

| Sample Depth (m) | Tissue Plot | Benthic Transect | Floe | Deep Sediment | Phytane ($\mu\text{g/g}$) | Pris/Phy | C ₁₈ /Phy | CPI | Status |
|------------------|-------------|------------------|-----------|---------------|-----------------------------|----------|----------------------|------------|-----------|
| 3 | 6 | | | | .05 | 1.0 | 2.0 | 2.5 | Oil |
| 3 | 8 | | -- | | .04 | 1.7 | .43 | 2.6 | Oil |
| 3 | 10 | | -- | | .27 | .68 | .70 | 1.4 | oil |
| 7 | 1 | -- | -- | | .01 | 5.9 | 1.2 | 2.8 | Low Oil |
| 7 | 5 | | -- | | .06 | 1.2 | .67 | 2.1 | Oil |
| 3 | | 1b | -- | -- | .18 | .67 | .39 | 1.8 | Oil |
| 3 | | 3b | -- | | .50 | .68 | .47 | 1.2 | Oil |
| 7 | -- | 1a | -- | -- | .04 | 2.7 | .92 | 2.3 | Oil |
| 7 | | 2C | -- | | .02 | 2.8 | 4.2 | 3.0 | oil |
| 7 | | 3b | -- | | .07 | 1.7 | .65 | 2.0 | Oil |
| 35 | | | | (4222) | .008 | 3.7 | 2.3 | 2.8 | Trace Oil |
| 35 | | | -- | (4221) | .010 | 3.0 | 3.7 | 3.0 | Low Oil |
| 3 | | -- | 6 | | .011 ^a | 1.4 | .81 | 2.4 | Low Oil |
| 3 | | | 8 | | .010^a | 2.4 | .90 | 1.7 | Low Oil |
| 3 | -- | | 10 | | .051^a | 0.9 | .22 | 1.7 | Oil |
| 7 | | -- | 1 | | .009^a | 11.0 | 3.0 | 1.6 | Trace Oil |
| 7 | | | 5 | | .008 ^a | 1.9 | 2.6 | 1.6 | Trace Oil |

-concentrations in $\mu\text{g/m}^2$.




FIGURE 3.7. GC² TRACES OF BAY 11 TISSUE PLOT SEDIMENT SATURATED HYDROCARBON* A- NO. 8; B- NO. 5.




FIGURE 3.8. AROMATIC PROFILES BY GC²/MS OF BAY 11 SEDIMENT


(.33, .90) mg/m^2 , at at 7 m, .23 (.17, .29) mg/m^2). Thus the 1983 values are very similar to those previously reported in 1982, in spite of the much larger bulk sediment (O-2 cm) reported in 1983. Thus, it appears that oil is being mixed into the upper sediment column in Bay 11 leaving a “steady state” concentration of oil in the floe layer.

3.2.1.2b Oil Composition by GC². A series of five Bay 11 floe samples were analyzed by GC2 (Table 3.4). Small to moderate quantities of oil were noted (Figure 3.9) in these samples when the petrogenic components (i.e. UCM material, phytane, etc.) were viewed against the background biogenic material. The saturated hydrocarbon weathering ratios (SHWR) of all of these samples approximated 1.0 indicating that the oil residues had been depleted of the lower molecular weight *n*-alkanes (C₁₀ to C₁₇). Petrogenic material in the floe at 3 meters depth was more highly biodegraded than that at 7m as judged by the C₁₈/Phy ratio which ranged from 0.2 to 0.9 at 3 m, with the lowest value corresponding to the highest oil concentration found in tissue plot #10. The C₁₈/Phy ratio was 2.5 to 3.0 at 7m depth.

3.2.1.2c Aromatic Hydrocarbon Composition by GC²/MS. The aromatic hydrocarbon composition of the sediment floe from Bay 11 was determined on three samples. The results are summarized in Figure 3.10. Aromatic compositions and concentrations were quite similar to those reported in the 1982 samples. For example, C₂phenanthrene and C₂ DBT were each present at - 300ng/m² in 1982. Comparable levels in 1983 were ~200ng/m².

3.2.1.3 Benthic Transects

3.2.1.3a Oil Concentrations by UV/F. Concentrations of oil in the sediments taken from the three transects are presented in Figure 3.6 and are summarized in Table 3.3. Concentrations in Transect 1 were similar at 3 and 7 m, 21.2 (8.6, 51.9) $\mu\text{g/g}$ and 22.5 (18.7, 27.1) $\mu\text{g/g}$ respectively, although the range (10.8-58.3 $\mu\text{g/g}$) was greater at the 3 m stations. Concentrations in Transect 2 were also similar at 3 m, 14.0 (4.7, 41.3) $\mu\text{g/g}$, and 7 m, 12.0 (10.9, 13.2) $\mu\text{g/g}$. Again the range at 3 m (5.3 to 45.1 $\mu\text{g/g}$) was greater owing to the patchiness observed repeatedly at the 3 m stations. Levels of oil in Transect 3 were much greater at 3 m, 119 (86.7, 162) $\mu\text{g/g}$ than at 7 m, 18.8 (9.9, 35.8) $\mu\text{g/g}$. This observation is both consistent with the observations for the adjacent tissue plots as well as with the 1982 observations.






FIGURE 3.10. AROMATIC PROFILES BY GC²/MS OF SEDIMENT FLOC FROM BAY 11

The 1983 oil levels in the **benthic** transects were much higher (4-10 times) than previously observed in 1982.

3.2. 1.3b Oil Composition by GC². Five **benthic** transect sediment samples were analyzed by GC² (Table 3.4). The compositional profiles of all of these samples closely resemble those from the tissue plots (see Figure 3.7). GC² analyses confirmed the presence of oil in **all** of these benthic transect samples. At lower concentrations the **PRIS/Phy** ratio became larger than 2.5 indicating an increased relative dominance of the **biogenic** isoprenoid, pristane. That the oil was moderately biodegraded is illustrated by the lower **C₁₈/Phy** ratio (0.4 to 1.0) compared to that in the undegraded oil. This extent of biodegradation is greater than was observed in samples taken from these locations in 1982. At that time, the lower extent of **C₁₈/Phy** ratio was ~1.0. Here in 1983, it has decreased further to 0.4 in some samples. The **CPI** values are all less than 3.0, indicating a higher degree of oiling than was the case in 1982, a finding that confirms the quantitative results presented previously.

3.2. 1.3c Aromatic hydrocarbon composition by GC²/MS. The results on the single benthic transect sediment sample analyzed by GC²/MS is presented in Figure 3. 10D. Low levels (5-10 ng) of individual aromatics were detected.

3.2.1.4 Microbiology Transects

3.2. 1.4a Oil Concentration and Composition by GC². A summary of the petroleum concentration data in the one microbiology transect is summarized along with relevant compositional parameters in Table 3.5. The locations of the transect is illustrated in Figures 3.11 and 3.12. Estimated petroleum concentrations ranged from 0.8 to 410 $\mu\text{g/g}$. Note that the total hydrocarbon levels are often higher at the lower levels than the estimated petroleum concentrations due to the presence of significant amounts, 2-3 $\mu\text{g/g}$, of biogenic hydrocarbons in petroleum-free sediment. Maximum concentrations were found at Station 14 and 13 at 2.4 and 4.0 meters depth, respectively. Concentrations decrease further offshore although petroleum residues were detected in all samples.

Representative compositional profiles are shown in Figures 3.13 and 3.14 and compositional parameters tabulated in Table 3.5. Note that a large amount of phytane present in these samples ($> .001$, background) directly indicates that large amounts of petroleum are present. The petroleum residues are moderately biodegraded with levels of

TABLE 3.5 SUMMARY OF MICROBIOLOGY SEDIMENT SAMPLE HYDROCARBON DATA.

| Bay | Water Depth (m) | Station | Total Hydrocarbon Concentration | Phytane (µg/g) | Pristane / Phytane | <u>n-C₁₈</u> / Phytane | C P I | Estimated Petroleum Concentration (µg/g) |
|-----|-----------------|---------|---------------------------------|----------------|--------------------|-----------------------------------|-------|--|
| 11 | 11.3 | 1 | 3.5 | .011 | 4.3 | .63 | 3.2 | 1.7 |
| 11 | 10.6 | 2 | 7.3 | .005 | 6.1 | 1.1 | 3.3 | .8 |
| 11 | 9.1 | 3 | 7.8 | .005 | 6.2 | 1.4 | 3.1 | .8 |
| 11 | 9.1 | 4 | 6.8 | .008 | 7.6 | 1.1 | 1.4 | 1.2 |
| 11 | 7.6 | 5 | 5.2 | .011 | 2.9 | .83 | 1.4 | 1.7 |
| 11 | 6.9 | 6 | 5.2 | .006 | 5.3 | 1.3 | 5.0 | .9 |
| 11 | 6.4 | 7 | 7.8 | .029 | 2.5 | .59 | 2.0 | 4.4 |
| 11 | 6.1 | 8 | 13 | .030 | 1.6 | .59 | 2.6 | 4.50 |
| 11 | 3.5 | 9 | 34 | .19 | .81 | .37 | 1.9 | 29.0 |
| 11 | 4.6 | 10a | 37 | .28 | .83 | .40 | 1.5 | 42.0 |
| | | 10b | 39 | .30 | .90 | .37 | 1.5 | 45.0 |
| | | 10C | 30 | .23 | .82 | .36 | 1.9 | 35.0 |
| 11 | 4.6 | 11 | 30 | .28 | 2.1 | .55 | 1.0 | 42.0 |
| 11 | 4.5 | 12 | 30 | .24 | 0.7 | .45 | 1.7 | 36.0 |
| 11 | 4.0 | 13 | 65 | .78 | .72 | .38 | 0.6 | 120.0 |
| 11 | 2.4 | 14 | 300 | 2.7 | .90 | .91 | 1.2 | 410.0 |
| 11 | 1.5 | 15 | 24 | .29 | .67 | .28 | 1.2 | 44 |
| 11 | 1.3 | 16 | 63 | 1.7 | .77 | .50 | 1.3 | 87.0 |
| 7 | 2 | 5 | 3.2 | .001 | 45 | 1.7 | 3.3 | <.5 |
| 7 | 4 | 4 | 3.7 | <.001 | | 4.0 | 7.4 | <.5 |
| 7 | 6 | 3 | 1.8 | <.001 | 33 | | 6.3 | <.5 |
| 7 | 8 | 2 | 1.9 | <.001 | | | 7.0 | <.5 |
| 7 | 10 | 1 | 5.1 | .012 | 3.1 | .45 | 1.7 | 1.8 |

^aIncludes any petroleum material plus biogenic compounds quantified by gas chromatography.

^bEstimated from known Phytane Content of Aged Lagomedio crude oil (6.4 mg Phytane/g oil) (from Boehm et al., 1982a).




FIGURE 3. 1. LOCATION OF BAY 11 MICROBIOLOGY SUBTIDAL SEDIMENT TRANSECT.




FIGURE 3.12. CROSS-SECTIONAL DEPTH PROFILE ALONG BAY 11 MICROBIOLOGY SEDIMENT TRANSECT




FIGURE 3.13. GC² TRACES IN MICROBIOLOGY SEDIMENT SAMPLES (BAY 11): A- STATION 12, 36 ppm (SATURATES); B- STATION 14, 410 ppm (SATURATES); C- STATION 14, 410 ppm (AROMATICS).

FIGURE 3.14. GC2 TRACES IN MICROBIOLOGY SEDIMENT SAMPLES (LOW LEVEL OIL): A- STATION 2, 0.8 ppm (SATURATES); B- STATION 5, 1.7 ppm (SATURATES).

n-C₁₈ substantially less than phytane in moderately to heavily oiled samples. A lesser degree of biodegradation is seen in the lightly oiled samples (~**1.0** ppm). Normal **alkanes** in the **C₁₀** to **C₁₇** range are quite prominent in the most heavily oiled sample (Station 14; Figure 3.1 3), thus resulting in a **SHWR** value of ~**2.0**. At **lower absolute** quantities of oil, the **SHWR** is lower 1.0- 1.5 indicating a higher degree of physical/chemical weathering. Stations 2 and 5 (Figure 3.14) contain small quantities of oil, computed from the Phytane/total oil ratio of 6.4 **mg/g**, which are obscured by the much larger amounts of **terrigenous** odd chain **n-alkanes**. Nevertheless, weathered oil is definitely present as noted by the presence of UCM feature on the GC² traces.

3.2.1.5 Sediment Cores

A series of four sediment cores, segmented in 5 cm segments to 15 cm depth in the sediment column were analyzed to examine vertical oil penetration in the sediment column.

3.2.1.5a Oil Composition by GC². Relevant compositional data from the saturated hydrocarbon GC² traces are summarized in Table 3.6. Concentrations of phytane are converted to "**total** estimated petroleum" by multiplying by 156 (the ratio of total oil to phytane in the Lagomedio oil). Very little oil penetration into the sediment is observed in these results. **Note** that the surface sediment sample is a 5 cm segment as compared with the normal 2 cm surface sediment sample obtained in this study. It can be therefore concluded that oil present in the Bay 11 sediments resides primarily in the top O-2 cm. We saw previously (Section 3.2.3.2a and 3.2. 1.2a) that oil **levels** had increased in surface sediment samples between 1982 and 1983, yet the surface floe levels remained similar. Oil, therefore, must have been mixed into the sediment to achieve a higher **bulk** sediment (O-2 cm) level without increasing floe values. Apparently, from the core data this penetration is superficial, not extending below the O-5 cm segment and undoubtedly residing **mostly** in the O-2 cm segment. This accounts for the low oil values shown by the core samples. Another explanation is that the differing sampling techniques biased the results, a likely contributing factor.

In any event, representative GC² traces an oiled core segment (3.1 **µg/g**) is shown in Figure 3.15 along with the companion 5-10 segment and the aromatic fraction from this latter segment. This **aromatic/olefenic** profile is typical of all of the sediment

TABLE 3.6 BAY 11 SEDIMENT CORE GC2 DATA

| Core | Section (cm) | Phy ($\mu\text{g/g}$) | $\frac{\text{Pris}}{\text{Phy}}$ | C₁₈ Phy | Estimated Oil Concentrations ($\mu\text{g/g}$) | Status |
|---------------------------|-----------------|----------------------------|----------------------------------|------------------------------|--|-----------|
| 11 N (3 meters) | 0-5 | .005 | >10 | 2.0 | 0.8 | Trace Oil |
| | 5-10 | .002 | 3.4 | 2.3 | <0.5 | --- |
| | 10-15 | .002 | 5.2 | 2.9 | <0.5 | --- |
| 11 s (3 meters) | 0-5 | .02 | .88 | .77 | 3.1 | oil |
| | 5-10 | .002 | 3.4 | 2.7 | <0.5 | --- |
| 11 N (7 meters) | 0-5 | .007 | 7.1 | 3.3 | 1.1 | oil |
| | 5-10 | .005 | 7.7 | 3.1 | .8 | Trace Oil |
| | 10-15 | <.001 | 55 | 17 | <0.5 | --- |
| 11 s (7 meters) | 0-5 | <.001 | 33 | -- | <0.5 | --- |
| | 5-10 | <.001 | 10 | -- | <0.5 | --- |

a - Estimated from **phytane/oil** ratio.




FIGURE 3.15. GC² TRACES OF BAY 11, S, 3m, SEDIMENT CORE: A- SATURATES 0-5 cm; B- SATURATES 5-10 cm; C- AROMATICS/UNSATURATES 5-10 cm; *- CYCLOALKENES.

samples. The aromatic fraction (Figure 3.15c) consists primarily of uncharacterized unsaturated hydrocarbon compounds of a biogenic or diagenetic origin. The saturate fraction of the 5-10 cm segment contains an abundance of **cycloalkenes** previously studied in other regions by Requejo and Quinn (1983). The Bay 11 north core at 3m illustrates no **visible** oil and only biogenic components (Figure 3.16).

3.2. 1.5b Aromatic Hydrocarbon Composition by GC²/MS. GC²/MS analytical results for two cores are presented in Figures 3.17 and 3.18. The 0-5 cm segment of the 11 south (3m) core contains individual petrogenic aromatics in the 2-10 rig/g range. Small quantities of the **alkylated** aromatics are also present at 5-10 cm. The entire 11 north (3m) core, Figure 3.18, contains only traces of the C₀ and C₁ phenanthrene compounds, 1-5 rig/g, which are typical **pre-spill** values. Very low levels of the C₀ and C₁ DBT compounds (probably not associated with oil) are seen in the 10-15 cm segment of the 11 north core. It is not clear why the 0-5 cm segment is essentially “free” of petrogenic aromatics, in spite of the clear evidence for the presence of these compounds in the 0-2 cm segment (i.e. the tissue plots).

3.2.1.6 Deep Sediments (35 m)

A series of four deep sediments taken in the offshore basin in the central Bay 11/ 12 area at 35m water depth were analyzed to determine longer range offshore transport of sedimented oil.

3.2.1.6a Oil Concentrations by UV/F. As the results in Tables 3.3 and 3.7 and Figure 3.19 indicate, petroleum residues were clearly detected by UV/F in these deep water sediments with concentrations ranging from 1.7 to 8.2 µg/g. It appears that oil has been transported to deeper areas offshore although the distributions are quite patchy.

3.2. 1.6b Oil Composition by GC². Two samples were analyzed by GC². Low levels of phytane (.008 and .01 µg/g) convert to 1.2 and 1.6 µg/g of oil respectively by GC². The GC² trace shown in Figure 3.20 shows the presence of C₁₅-C₂₀ petroleum alkanes along with phytane and a small UCM, all characteristic of petroleum inputs to the deep sediments.

3.2. 1.6c Aromatic Hydrocarbon Composition by GC²/MS. The one deepwater sediment sample examined (B4221) contained 1-3 rig/g of the C₀ and C₁ phenanthrene compounds but not detectable petroleum aromatics in spite of the clear UV/F signal of the total extract which yielded a result of 8.2 µg/g of oil.




FIGURE 3.16. GC² TRACES OF SATURATED HYDROCARBONS IN BAY 11 N, 3m, SEDIMENT **CORE:** A- 0-5 cm; B- 5-10 cm; C- 10-15 cm.




FIGURE 3.17. AROMATIC PROFILES BY GC²/MS OF BAY 11 SEDIMENT CORE 11S, 3m




FIGURE 3.18. AROMATIC PROFILES BY GC²/MS OF BAY 11 SEDIMENT CORE 11N, 3m

TABLE 3.7 BAY 11/12 DEEP WATER DREDGES SEDIMENT PETROLEUM
HYDROCARBONS BY UV/F

| Sample I.D. | Petroleum Hydrocarbons ($\mu\text{g/g}$) |
|--------------|--|
| B4221 | 8.2 |
| B4222 | 5.5 |
| B4223 | 4.4 |
| B4224 | 5.9 |
| B4225 | 1.7 |




FIGURE 3.19. UV/F SPECTRA OF BAY 11/12 DEEPWATER SEDIMENT SAMPLE.




FIGURE 3.20. GC2TRACE OF DEEPWATER SEDIMENT SAMPLE;
SATURATED HYDROCARBONS

3.2.1.7 Bay 11 Beach Sediments

The concentrations of oil found in beach samples from Bay 11 along with details of the composition of this oil are presented in Table 3.8. These oil residues represent the source of material for the Bay 11 subtidal sediments and perhaps to other sediments in the area. Note that the total extractable petroleum values include polar (i.e. non-hydrocarbon) material as well as petroleum hydrocarbons.

Concentrations of oil on the Bay 11 beach surface were quite patchy ranging from 72 to 19,400 $\mu\text{g/g}$. The mean concentration, with lower and upper 95% confidence limit values (log transformed data) was 1250 (192, 8090) $\mu\text{g/g}$. Sub-surface oil values were lower on the average 158 (5.9, 4220) $\mu\text{g/g}$ although the range is very large.

Compositional details are also presented in Table 3.8 and two representative GC2 traces are shown in Figure 3.21.

A range of weathering states are observed. The X 1 surface sample exhibits an SHWR = 2.0 and, ALK/ISO value of 2.2 and an AWR of 2.5 all indicative of lightly weathered oil. The GC² trace of this sample is shown in Figure 3.2 1a. At the other compositional extreme (Figure 3.20b) observed for these samples in the Profile 6, lower surface sample (SHWR = 1.0; ALK/ISO = 0.3). The oil in this sample is highly weathered from both physical/chemical and microbial degradation viewpoints. In general, the lower surface samples are much more highly weathered than their counterparts on the upper end of the beach profile. Many intermediate values are observed.

3.2.2 BAY 9

Three types of sediment samples were analyzed from the Bay 9 beach and **subtidal** region during 1983. Sediment samples (0-2 cm) were analyzed from the 7m tissue plots and from the 3m benthic transect stations. In addition, a series of seven Bay 9 beach samples were analyzed.

3.2.2.1 Tissue Plots

3.2.2.1a Oil Concentrations by UV/F. The concentrations of oil in the Bay 9 **subtidal** sediments are presented in Figure 3.22. Concentrations of oil in the 7m tissue

TABLE 3.8 SHORELINE STUDY 1983 HYDROCARBON CHEMISTRY RESULTS - ANALYTICAL RESULTS; RAGGED CHANNEL BEACHES.

| Sample ID | Plot | Depth | Saturated Hydrocarbon µg/g ^a | Aromatic Hydrocarbon µg/g ^a | Total Petroleum Hydrocarbons µg/g ^a | Total Extractable Organics ^b µg/g | SHWR | ALK/ISO | AWR |
|-----------|-----------------|------------------------|---|--|---|---|------|---------|-----|
| 4124 | Bay 9 | 100, Upper Surface | 0.0 | 1.0 | 1.0 | 8.1 | 2.0 | 0.8 | 1.2 |
| 4126 | Bay 9 | 100, Mid-Surface | 0.9 | 0.4 | 1.3 | 6.9 | 1.2 | 1.4 | NA |
| 4127 | Bay 9 | 100, Mid Sub-Surface | 0.0 | 0.0 | 0.0 | 11.3 | 1.7 | 0.9 | 1.6 |
| 4128 | Bay 9 | 100, Lower Surface | 0.2 | 0.0 | 0.2 | 2.92 | ND | ND | NA |
| 4129 | Bay 9 | 100, Lower Sub-Surface | 0.8 | 0.8 | 1.6 | 11.6 | 1.0 | 0.9 | NA |
| 4952 | Bay 9 | 300, Upper Surface | 0.7 | 0.2 | 0.9 | 10.8 | 1.1 | 1.5 | NA |
| 4130 | Bay 9 | 300, Mid-Surface | 0.5 | 1.1 | 1.6 | 6.5 | 1.2 | 0.8 | NA |
| 4132 | Bay 9 | 300, Lower Surface | 0.0 | 0.7 | 0.7 | 9.7 | 1.2 | 1.6 | NA |
| 4134 | Bay 11 | 2, Upper Surface | 221. | 180. | 401. | 1,260. | 1.0 | 1.0 | 1.0 |
| 4135 | Bay 11 | 6, Upper Surface | 8,490. | 5,380. | 13,900. | 17,300. | 1.6 | 1.9 | 2.2 |
| 4136 | Bay 11 | 2, Mid-Surface | 601. | 307. | 908. | 1,880. | 1.1 | 0.4 | 1.1 |
| 4137 | Bay 11 | 6, Mid-Surface | 9,210. | 4,280. | 13,500. | 16,700. | 1.7 | 2.1 | NA |
| 4138 | Bay 11 | 2, Lower Surface | 55.1 | 16.9 | 72.0 | 197. | 1.1 | 0.2 | NA |
| 4139 | Bay 11 | 6, Lower Surface | 2,580. | 1,310. | 3,890. | 6,160. | 1.0 | 0.3 | NA |
| 4140 | Bay 11 | 4, Upper Surface | 12,200. | 7,230. | 19,400. | 25,400. | 1.6 | 2.5 | NA |
| 4141 | Bay 11 | 8, Upper Surface | 2,380. | 1,100. | 3,480. | 4,810. | 1.1 | 1.4 | NA |
| 4142 | Bay 11 | 4, Mid-Surface | 4,220. | 1,930. | 6,150. | 10,800. | 1.0 | 0.7 | NA |
| 4143 | Bay 11 | 8, Mid-Surface | 1,010. | 503. | 1,510. | 2,450. | 1.0 | 1.4 | NA |
| 4144 | Bay 11 | 4, Lower Surface | 29.6 | 13.1 | 42.7 | 94.0 | 1.2 | 0.4 | NA |
| 4145 | Bay 11 | 8, Lower Surface | 835. | 422. | 1,260. | 2,000. | 1.1 | 0.3 | NA |
| 4146 | Bay 11 | X1 Surface | 16,200. | 1,050. | 17,200. | 26,900. | 2.0 | 2.2 | 2.5 |
| 4147 | Bay 11 | X2 Surface | 216. | 107. | 322. | 600. | 1.3 | 1.2 | NA |
| 4148 | Bay 11 | X3 Surface | 332. | 186. | 518. | 809. | 1.8 | 2.1 | NA |
| 4149 | Bay 11 | X4 Surface | 122. | 69.8 | 192. | 730. | 1.6 | 1.8 | 2.1 |
| 4150 | Bay 11 | X5 Surface | 7,670. | 5,130. | 12,800. | 15,900. | 1.4 | 1.0 | NA |
| 4946 | Bay 11 | X1 Sub-Surface | 4,640. | 2,380. | 7,020. | 8,980. | 1.6 | 2.0 | 2.3 |
| 4947 | Bay 11 | X2 Sub-Surface | 41.4 | 18.6 | 78.6 | 151.0 | 1.2 | 0.7 | NA |
| 4948 | Bay 11 | X3 Sub-Surface | 2.4 | 2.3 | 4.7 | 72.2 | 1.3 | 1.4 | NA |
| 4949 | Bay 11 | X4 Sub-Surface | 0.9 | 3.0 | 3.9 | 227. | 1.3 | 1.4 | 3.5 |
| 4950 | Bay 11 | X5 Sub-Surface | 3,430. | 2,180. | 5,600. | 7,180. | 1.8 | 1.8 | NA |
| 4951 | Bay 11 | X6 Sub-Surface | 164. | 105. | 269. | 688. | 1.7 | 2.1 | NA |
| 4955 | Crude Oil Point | X7 Surface | 1,060. | 467. | 1,530. | 2,600. | 1.0 | 1.2 | NA |
| 4956 | Crude Oil Point | X8 Sub-Surface | 1,220. | 724. | 1,940. | 3,000. | 1.0 | 1.6 | NA |

a .determined gravimetrically.

b . contain petrogenic hydrocarbons, petrogenic polar compounds and biogenic compounds.

ND. none detected.




FIGURE 3.21. GC²TRACE OF BAY 11 BEACH SEDIMENTS: A- SATURATES S4146 (UNWEATHERED, **UNDEGRADED**); B- SATURATES S4139 (WEATHERED AND DEGRADED); *-UNIDENTIFIED CYCLIC ALKANES (TERPENOID).




FIGURE 3.22. BAY 9 SEDIMENT PETROLEUM HYDROCARBON CONTENT; BY UV/F (AUGUST 15, 1983).
(NA = NOT ANALYZED)

plots ranged from 5.6 to 9.5 $\mu\text{g/g}$ with a mean of 7.6 (6.0, 9.5) $\mu\text{g/g}$ as determined by UV/F. These values are higher than those reported for the 1982 samples which were 2.2 (1.5, 3.1) $\mu\text{g/g}$.

3.2.2. 1b Oil Composition by GC2. Three tissue plot sediments from Bay 9 were analyzed by GC² to confirm the presence of oil and to examine the sample's hydrocarbon composition. As can be seen from one of the GC² traces (Figure 3.23) low levels of identifiable petrogenic compounds (n-alkanes C₁₃-C₁₉, and phytane) are seen in the samples. Phytane is detected at levels of .002 to .005 $\mu\text{g/g}$ which convert to concentrations of 0.3 to 0.8 $\mu\text{g/g}$ compared with the much higher levels detected by UV/F. These discrepancies are examined in Section 3.2.5. The aromatic/olefinic fraction exhibits large amounts of biogenic/diagenetic unsaturates with possibly minor quantities of petroleum aromatics. These aromatics are discussed in the next section.

3.2.2. 1c Aromatic Hydrocarbons Compositions by GC²/MS. Two Bay 9 a tissue plot samples (B4054, 7m, No. 5 and B4077, 7m, No. 3) analyzed by GC²/MS contained low levels 1-3 $\mu\text{g/g}$ of petroleum aromatics (i.e. the alkylated phenanthrene and DBT compounds. These values were a factor of 5 lower than the levels observed in 1982. Note however, that the levels determined on the 1983 samples are quite similar to those reported after the spills in 1981 when bulk oil levels in the Bay 9 sediments were roughly the same concentration (1981: mean = 9.0 $\mu\text{g/g}$).

3.2.2.2 Benthic Transects

3.2.2.2a Oil Concentrations by UV/F. Oil concentrations in the 3m benthic transects as determined by UV/F were as follows: Transect 1 = 1.0 (.62, 1.8) $\mu\text{g/g}$; Transect 2 = 4.6 (3.3, 6.4) $\mu\text{g/g}$; Transect 3 = 5.2 (4.4, 6.1) $\mu\text{g/g}$. The values in Transects 2 and 3 are higher than those reported on 1982 field samples at which time Transect 2 levels were .52 (.23, 1.2) $\mu\text{g/g}$ and Transect 3 levels were .91 (.83, 1.5) $\mu\text{g/g}$. Levels in Transect 1 were similar in 1982 and 1983 (1982 Transect 1 = 1.0 (.66, 1.6) $\mu\text{g/g}$). Thus it appears that levels of 3-7 $\mu\text{g/g}$ of oil were added to the benthic transect samples. These levels approximate those observed in the 1982 Bay 9 microbiology samples 3-6 $\mu\text{g/g}$ in 1982.

3.2.2.2b Oil Composition by GC2. GC² analyses of the two Bay 9 benthic transect samples analyzed reveal the presence of only low levels of any petrogenic compounds in the saturated hydrocarbon fraction (similar to Figure 3.23). Levels of

FIGURE 3.23. GC² TRACE OF BAY 9 TISSUE PLOT **SEDIMENT**: **A**- SATURATED HYDROCARBONS; **B**- AROMATIC/UNSATURATED HYDROCARBONS.

phytane in the two samples analyzed were detectable, but very low (.005 and .004 $\mu\text{g/g}$) albeit higher than the .001 ($\mu\text{g/g}$) background. However, these phytane levels convert to oil levels of .8 and .6 $\mu\text{g/g}$ respectively, much lower than the UV/F determined values.

3.2.2.2c Aromatic Hydrocarbon Composition by GC²/MS. The single benthic transect sample analyzed by GC²/MS (B4088; 3m; Benthic Station 3a) contained individual aromatic hydrocarbon levels of 1-3 rig/g (alkylated phenanthrene and DBT compounds) amidst a 3 rig/g background of polycyclic aromatic hydrocarbons (4 and 5 rings).

3.2.2.3 Bay 9 Beach

The analytical results for a set of eight Bay 9 beach sediment samples are presented in Table 3.8. These oil concentrations range from 0.5 to 1.6 $\mu\text{g/g}$, in the range of those levels reported in 1982. The highest level samples were comprised of a weathered (SHWR = 1.2) undegraded (C₁₈/Phy = 3.4) saturated hydrocarbon composition (Figure 3.24). This oil is significantly different from that observed on the Bay 11 beach, as it is undegraded.

3.2.3 BAY 7

Two types of sediment samples were analyzed from the Bay 7 subtidal area: five tissue plot sediments and nine benthic transect samples.

3.2.3.1 Tissue Plots

3.2.3.1a Oil Concentrations by UV/F. UV/F-determined values of oil concentrations in Bay 7 stations (Figure 3.25) averaged 3.2 (1.4, 7.4) $\mu\text{g/g}$. Aside from the one high value, 12.8 $\mu\text{g/g}$ / tissue plot # 4, the mean value is 2.3 (1.6, 3.3) $\mu\text{g/g}$, approximately twice the mean value determined in 1982, 1.2 (.96, 1.4) $\mu\text{g/g}$.

3.2.3.1b Oil Composition by GC². The existence of petroleum at tissue plot #4 was confirmed by the GC² data on this sample. This sample, shown in Figure 3.26A, contains .05 $\mu\text{g/g}$ phytane (~7.8 $\mu\text{g/g}$ of oil) and a small, but significant quantity of UCM material. The C₁₈/Phy ratio of 0.5 indicates that this petroleum material is significantly biodegraded. This ratio is equal to 1.6 in undegraded oil. The PRIS/Phy ratio of 2.8




FIGURE 3.24. GC² TRACE OF REPRESENTATIVE BAY 9 BEACH SEDIMENT; SATURATED HYDROCARBONS.




FIGURE 3.25. BAY 7 SEDIMENT PETROLEUM HYDROCARBON CONTENT; BY UV/F (AUGUST 14, 1983).
(NA = NOT ANALYZED)




FIGURE 3.26. GC² TRACE BAY 7 SEDIMENTS: A- SATURATES TISSUE PLOT NO. 4; & SATURATES **BENTHIC** TRANSECT.

indicates that **biogenic** pristane “overprints” any **petrogenic** pristane (**PRIS/PHY** in oil = 0.74). **Petrogenic phytane was detected in tissue plot # 5 (.008 µg/g),** tissue plot #3(.009) and the agreement between UV/F and GC² results in Bay 7 is good (see Section 3.2.5).

3.2.3.1c Aromatic Hydrocarbon Composition by GC²/MS. Two samples (Plot No. 4a and 5) were analyzed by **GC²/MS.** The presence of petroleum in plot 4, corresponding to the Figure 3.26A sample, was confirmed (Figure 3.27A). Individual aromatics were present at the 1-4 rig/g level and a petroleum composition was noted, as opposed to the background distribution (Figure **3.27B**) observed for the other sample from Bay 7.

3.2.3.2 Benthic Transects

3.2.3.2a Oil Concentrations by UV/F. Concentration data for the Bay 7 **benthic** transect sediments is shown in Figure 3.25. Values are low as follows: Transect 1 = 2.1 (1.9, 2.4) **µg/g;** Transect 2 = 1.5 (.99, 2.4) **µg/g;** Transect 3 = 1.0 (.55, 1.8) **µg/g.** These values are similar to those observed in the tissue **plots** in Bay 7 in 1982. No 3m samples were analyzed in 1982.


3.2.3.2b Oil Composition by GC². Two samples were analyzed to examine hydrocarbon compositions of the Bay 7 **benthic** transects. One of the two saturated hydrocarbon GC² traces is presented in Figure **3.26B.** No phytane and hence no petroleum residues were detected in these 3m **samples.**

3.2.4 MILNE INLET SAMPLES

Five sediment samples taken from the **subtidal** area of **Milne** Inlet on the western side of Ragged **Island** were analyzed to determine if any oil residues were detectable in these reference areas.

3.2.4.1 Oil Concentrations by UV/F

No oil was detected in any of the five samples UV/F spectra (e.g., Figure 3.28) contained only small 350 nm responses. Resultant oil concentrations are shown in Table 3.9. The mean value is 0.78 **µg/g,** essentially a background **level.**

FIGURE 3.27. AROMATIC PROFILES BY **GC²/MS** OF BAY 7 SEDIMENTS




FIGURE 3.28. UV/F SPECTRA OF **MILNE** INLET SEDIMENT SAMPLE DILUTION SERIES, SHOWING NO OIL PRESENT.

TABLE 3.9 MILNE INLET SUBTIDAL SEDIMENT CONTROL SAMPLES -
PETROLEUM HYDROCARBON CONTENT BY **UV/F**

| Sample I.D. | Plot | Petroleum Hydrocarbons ($\mu\text{g/g}$) |
|--------------|------|--|
| B4091 | 1 | 0.8 |
| B4092 | 2 | 0.7 |
| B4093 | 3 | 1.1 |
| B4094 | 4 | 0.5 |
| B4095 | 5 | 0.8 |

3.2.4.2 Oil Compositions by GC²

The two GC 2 analyses performed on the Milne Inlet samples confirm the purely biogenic composition of these samples. No phytane or UCM was detected in either sample. The composition consisted of terrigenous **n-alkanes** and **olefinic** material (Figure 3.29).

3.2.4.3 Aromatic Hydrocarbon Composition by GC²/MS

The aromatic hydrocarbon composition of the one Milne Inlet sample examined is illustrated in Figure 3.30.

3.2.5 COMPARISON OF UV/F-DERIVED PETROLEUM CONCENTRATIONS AND GC²-DERIVED RESULTS

A large discrepancy exists between UV/F and **GC²-derived** hydrocarbon values as shown in Table 3.10. This discrepancy was noted in the 1982 study results (Boehm, 1983a) and has become wider with time. The original close agreement between these two sets of results Boehm et al. (1982a) apparently does not hold as oil residues get progressively more weathered. As we compute a **GC²-derived** value from the phytane concentrations (i.e. **Phytane/total oil** = 6.4 mg/g phytane oil) this computation becomes less reliable as phytane itself is degraded. We know that the UV/F 350-360 nm band was nearly absent in **prespill** samples and is also absent in Milne Inlet sediment samples. Therefore, we must conclude that the UV/F data is reliable at least in a comparative sense when viewed against 1981-1982 data.

Note also that as the oil weathers, the correlation of the 350-360 nm intensity on the UV/F spectra with total oil concentrations breaks down. However, we conclude that the UV/F values should be used as the petroleum concentrations in this study.

3.2.6 REANALYSIS OF 1982 FIELD SAMPLES

As part of the QC/QA program performed as part of this study, three sediment samples from the 1982 field program representing a range of concentrations were




FIGURE 3.29. GC2 TRACE **MILNE** INLET SEDIMENT SATURATED HYDROCARBON, ● -CYCLOALKENES.




FIGURE 3.30. AROMATIC PROFILES BY **GC²/MSMILNE** INLET SEDIMENT SAMPLE

TABLE 3.10 **UV/F** VERSUS GC2 - DERIVED RESULTS ON SEDIMENTS

| Bay | Tissue Plot | Benthic Transect | Oil Concentration UV/F ($\mu\text{g/g}$) | Phytane ($\mu\text{g/g}$) | Oil Concentration GC2 ($\mu\text{g/g}$) |
|-----|-------------|------------------|--|--------------------------------|---|
| 11 | 1 | | 21.2 | .01 | 1.5 |
| | 5 | | 20.5 | .06 | 9.2 |
| | 6 | | 22.7 | .05 | 7.7 |
| | 8 | | 16.8 | .04 | 5.6 |
| | 10 | | 85.6 | .27 | 42 |
| | | 1b(3m) | 58.3 | .18 | 28.5 |
| | | 3b(3m) | 137 | .50 | 79.3 |
| | | 1a(7m) | 21.8 | .04 | 5.5 |
| | | 2c(7m) | 10.7 | .02 | 3.0 |
| | | 3b(7m) | 39.0 | .07 | 10.8 |
| 9 | 2 | | 7.4 | .002 | 0.3 |
| | 3 | | 9.6 | .003 | 0.5 |
| | 5 | | 9.5 | .003 | 0.5 |
| | | 2c(3m) | 6.9 | .005 | 0.8 |
| | | 3a(3m) | 6.2 | .004 | 0.6 |
| | 3 | | 1.6 | .008 | 1.3 |
| | 4 | | 12.8 | .05 | 7.8 |
| | 5 | | 1.8 | .009 | 1.4 |
| | | 2a(3m) | 2.5 | .002 | 0.3 |
| | | 3b(3m) | 1.6 | .003 | 0.5 |

a **Phytane** x **156** . Oil Concentration.

reanalyzed here. Other samples (e.g., tissues and shoreline sediments) were not available for this phase of the program. The UV/F and GC2 results are summarized in Table 3.11. The agreement between the two data sets is generally quite good.

3.3 Oil In Marine Organisms

3.3.1 Mya truncata

Mya samples were analyzed from each of the 7 meter tissue plots in Bays 11, 9 and 7. Following UV/F analyses the extracts from each bay were pooled and one combined sample was analyzed by GC² and GC²/MS; to determine compositional data.

3.3.1.1 Bay 11

3.3.1. 1a Oil Concentrations by UV/F. Levels of UV/F-determined petroleum in the Bay 11 Mya samples ranged from <1.0 µg/g to 10.6 µg/g (dry weight basis). The concentrations, summarized in Figure 3.31, 4.0 (1.7, 9.7) µg/g are similar to those reported on the September 1982 samples, 4.7 (4.0, 5.7) µg/g although the distributions are more patchy in 1983.

3.3.1. 1b Oil Composition by GC². The composition of saturated hydrocarbon fraction in Bay 11 Mya is shown by the GC² trace in Figure 3.32a. The three major features of the GC² trace are:

1. The significant quantity of UCM material indicative of weathered petroleum,
2. The terrigenous n-alkane content indicating that the animals contained sediment material,
3. The prominence of the isoprenoid hydrocarbons which indicate the oil in the animals was biodegraded.

Note that the GC² trace was expanded vertically to give the Figure 3.32A illustration.

The C₁₈/Phy ratio is 1.0 in this composite sample. This suggests that the animals are still acquiring oil from the Bay 11 system because in previous years this ratio




TABLE 3.11. QC/QA ANALYSES: COMPARISON OF RESULTS IN 1982 FIELD SAMPLES

| | Oil Concentrations by UV/F ($\mu\text{g/g}$) | | Phytane ($\mu\text{g/g}$) | | Oil Concentration by Phytane Conversion ($\mu\text{g/g}$) | | Pris/Phy | | C ₁₈ /Phy | |
|----------------------------|--|----------|--------------------------------|----------|--|----------|----------|----------|----------------------|----------|
| | <u>a</u> | <u>b</u> | <u>a</u> | <u>b</u> | <u>a</u> | <u>b</u> | <u>a</u> | <u>b</u> | <u>a</u> | <u>b</u> |
| 11 Plot No. 10 (187) | 66 | 84 | .41 | .60 | 64 | 94 | .87 | 1.3 | .91 | 1.3 |
| 11 Plot No. 5 (37) | 49 | 32 | .06 | .04 | 9.4 | 6.2 | 1.4 | 1.3 | 1.2 | 1.3 |
| 7 Plot | 2.2 | 2.1 | .003 | .005 | 0.5 | 0.8 | — | 10 | 5.2 | 30 |

1983 analyses (ERCO)

1984 analyses (BATTELLE)

TISSUE PLOTS



^aMean(Lower 95%, Upper 95%)

FIGURE 3.31. SUMMARY OF OIL CONCENTRATIONS IN Mya truncata, by UV/F, (µg/g dry wt.).

FIGURE 3.32. Mya truncata SATURATED HYDROCARBON GC² DETERMINATIONS.

had decreased to values much less than 1.0. The **PRIS/Phy** ratio in the animals **equals ~14** suggesting that the marine **biogenic** material is quantitatively more important than the **petrogenic** input, a fact borne out by the biogenic character of the GC2 trace in Figure 3.32a. Significant quantities of UCM material confirm the petroleum residues.

As a confirmatory step in the UV/F determinations, the combined f2 fractions from all five samples was **analyzed by UV/F**. The resultant **value** of 7.5 $\mu\text{g/g}$ is close to the arithmetic mean of the five separate plots (5.2 $\mu\text{g/g}$).

3.3.1.1c Aromatic Hydrocarbon Composition by GC²/MS. The composite aromatic hydrocarbon fraction of this sample contained only very low quantities of petroleum aromatics (Figure 3.33). **Alkylated** naphthalenes were present at the 1-5 rig/g level, down from the 10-15 rig/g levels observed in 1982. The **alkylated phenanthrenes** and dibenzothiophenes which were the dominant components in the 1982 animals are present only in trace levels 0.5 to 1.0 rig/g in the 1983 field samples. The **alkylated naphthalenes** are now dominating the aromatic composition albeit at very **low levels (2-5 rig/g)**. Comparable levels in the summer of 1982 were 1-15 rig/g of individual aromatic compounds.

3.3.1.2 Bay 9

3.3. 1.2a Oil Concentrations by UV/F. Mya samples from Bay 9 contain 2.9 (1.7, 5.2) $\mu\text{g/g}$ of petroleum by UV/F (Figure 3.31). These concentrations are higher than those reported for the 1982 field samples at which time levels were near or at background levels 0.81 (.52, 1.3) $\mu\text{g/g}$. Confirmatory UV/F analysis of a combined f2 fraction **yielded a value** of 3.3. $\mu\text{g/g}$ **very close to** the mean of the values determined on unfractionated extracts.

3.3.1.2b Oil Composition by GC². GC2 analysis of the combined sample yielded no definitive petrogenic component. A biogenic assemblage dominated with a **PRIS/Phy** value >50 and absolute phytane levels <0.001 $\mu\text{g/g}$.

3.3.1.2c Aromatic Hydrocarbon Composition by GC²/MS. GC²/MS analysis of the Bay 9 **Mya** sample yielded no detectable aromatic hydrocarbons (i.e. <1 rig/g). This represents a tenfold decrease in **identifiable** aromatics as the 1982 values were in the 2-10 rig/g range.




FIGURE 3.33. Mya AROMATIC PROFILES BY GC²/MS (BAY 11).

3.3.1.3 Bay 7

3.3.1.3a Oil Concentration by UV/F. UV/F determined levels in Mya from Bay 7 (Figure 3.31) were 1.1 (.83, 1.6) $\mu\text{g/g}$ which represents values near the detection limit (0.8 $\mu\text{g/g}$) and essentially containing only the smallest traces of a 355 nm shoulder on the UV/F trace. The combined f 2 UV/F value 1.1 $\mu\text{g/g}$ was identical to the mean of the individual samples.

3.3.1.3b Oil Composition by GC2. The saturated hydrocarbon GC2 trace exhibits a nearly petroleum-free composition (Figure 3.32B) except for a low quantity of UCM material and detectable phytane levels (-0.02 $\mu\text{g/g}$).

3.3.1.3c Aromatic Hydrocarbon Composition by GC²/MS. Bay 7 Mya samples contained only trace levels 1-2 $\mu\text{g/g}$ of alkylated naphthalenes and no phenanthrene or DBT compounds. When sampled in 1982, the comparable values were 2-8 $\mu\text{g/g}$.

3.3.2 Serripes groenlandicus

Samples of Serripes were collected and processed in a manner identical to that used for Mya (see Section 3.3.1)

3.3.2.1 Bay 11

3.3.2.1a Oil Concentrations by UV/F. The concentration summary of the Bay 11 animals is presented in Figure 3.34 and the UV/F trace of the combined f2 fractions shown in Figure 3.35. The concentrations in the tissue plots 10.9 (7.0, 17.0) $\mu\text{g/g}$ is verified by the combined f2 result, 9.1 $\mu\text{g/g}$. Oil levels in Bay 11 Serripes when previously sampled in 1982 were 5.2 (4.1, 6.4) $\mu\text{g/g}$. Thus the values in Serripes have increased by a factor of two on the average.

3.3.2.1b Oil Composition by GC². The GC² trace shown in Figure 3.36 exhibits low levels of petrogenic alkanes in the n-C₂₀ to n-C₃₀ range, a significant quantity of UCM material and the presence of .07 μg phytane, the latter of which converts to 11.2 $\mu\text{g/g}$ of petroleum. Thus the GC² results confirm the levels of oil determined by UV/F and indicate that the oil present is biodegraded in the n-C₁₀ to n-C₂₀ range with petrogenic alkanes still present in the samples.

TISSUE PLOTS

| | | | | | | |
|------------|-------------|-------------|------------|-------------|--------------|-----------------|
| 7.9 | 14.5 | 16.0 | 5.9 | 14.3 | BAY 11 7m | 10.9(7.0, 17.0) |
| 1 | 2 | 3 | 4 | 5 | | |

| | | | | | | |
|----------|----------|----------|----------|---------------|-------------|-----------------|
| <.8 | 2.8 | <.8 | <.8 | <.8 | BAY 9 7m | 1.0(,) |
| 1 | 2 | 3 | 4 | 5 | | |

| | | | | | | |
|----------|----------|----------|----------|------------|-------------|-------------------------------------|
| 3.1 | 0.9 | 0.8 | 23.4 | 0.8 | BAY 7 7m | 1.2(.59, 2.2) (Excluding Plot 4) |
| 1 | 2 | 3 | 4 | 5 | | |

^aMean(Lower 95%, Upper 95%)

FIGURE 3.34. **SUMMARY OF OIL CONCENTRATIONS IN Serripes groenlandicus, UV/F, (ug/g dry wt.).**





FIGURE 3.35. UV/F SPECTRA OF *Serripes* SAMPLE EXTRACT FROM BAY 1 I
COMBINED F2 FRACTION.



3.3.2.1c Aromatic Hydrocarbon Composition by GC²/MS. Analysis of the composite Serripes sample by GC²/MS, detected trace quantities of **alkylated naphthalenes** (3 rig/g) and phenanthrenes (2 rig/g) with no DBT compounds detected. This represents a drastic decrease from the 10-100 rig/g values determined in 1982 (Boehm 1983a).

3.3.2.2 Bay 9

3.3.2.2a Oil Concentrations by UV/F. One of the Bay 9 Serripes UV/F spectra is shown in Figure 3.37. These values if quantified blindly would indicate the presence of large quantities of oil in these animals and indeed the presence of oil is suggested by the UV/F traces (e.g., Figure 3.37). However, when the extract was fractionated and the f2 was rerun by UV/F, the 355 nm peak was removed from the sample. We suspect that a polar, non-petroleum interference having a spectral maximum at ~380 nm contributed to the spurious 355 nm “oil peak” in the unfractionated extracts. Essentially, oil was detected in only one sample (Plot 2). The actual mean value which should be at 2.8 µg/g.

3.3.2.2b Oil Composition by GC². The Bay 9 Serripes composite gave a GC² saturated hydrocarbon trace as shown in Figure 3. 36B. No detectable petroleum is seen in the sample. Only biogenic components were detected, confirming the absence of oil in Bay 9 Serripes.

3.3.2.2c Aromatic Hydrocarbons by GC²/MS. No petroleum aromatics were detected in the Bay 9 Serripes composite.

3.3.2.3 Bay 7

3.3.2.3a Oil Concentrations by UV/F. The summary of the Bay 7 Serripes oil concentrations (Figure 3.34) shows values ranging from <0.8 µg/g to 23.4. We suspect that as in the Bay 9 samples the 23.4 value may be a spurious result, due to the spectral interference identical for that in Bay 9 (Figure 3.37B). However, it is interesting to note that this Serripes oil value coincided with the highest sediment oil value of 12 ppm at tissue plot 4 in Bay 7. The mean of the other four samples is 1.2 µg/g which is lower than the 2.2 µg/g value reported on the 1982 samples.




FIGURE 3.37. Serripes UV/F SPECTRA FROM BAYS 9(A) and 7(B) SHOWING NON-PETROLEUM SPECTRAL INTERFERENCE WHICH WAS ELIMINATED BY COLUMN CHROMATOGRAPHY.

3.3.2.3b Oil Composition by GC². The Bay 7 GC² trace confirmed the absence of oil in these samples. Only **biogenic** material was detected, the hydrocarbon assemblage being dominated largely by pristane.

3.3.2.3c Aromatic Hydrocarbon Composition by GC²/MS. The Bay 7 Serripes sample contained no detectable petroleum aromatics.

3.3.3 Macoma calcaria

Samples **were** collected and processed as were Mya (Section 3.3.1).

3.3.3.1 Bay 11

3.3.3.1a Oil Concentrations by UV/F. The Bay 11 Macoma samples exhibited high concentrations of oil (Figure 3.38). The presence of oil was clearly discernible from the UV/F spectra (Figure 3.39). The concentrations in Bay 11, 63.8 (~~44.0~~, 92.7) µg/g are **nearly** identical to those determined from the 1982 samples 60.0 (39.0, 92.0) µg/g.

3.3.3.1b Oil Composition by GC². Moderate quantities of heavily weathered oil were detected in the combined Macoma samples including -0.6 µg/g phytane (=94 µg/g oil by GC²) and a significant amount of UCM material (Figure 3.40). The isoprenoids pristane and phytane are prominent confirming the highly biodegraded nature of the oil residues. This composition is similar to that observed in 1982. The smooth *n*-alkane (C₂₀-C₃₀) distribution further confirms the presence of considerable quantities of petroleum.

3.3.3.1c Aromatic Hydrocarbons Composition by GC²/MS. Only the alkylated phenanthrene compounds were detected in the composite Macoma sample (Figure 3.41). The highest value of 12 ng/g is over an order of magnitude lower than the comparable values determined in 1982.

3.3.3.2 Bay 9

3.3.3.2a Oil Concentrations by UV/F. Macoma samples from Bay 9 contained 12.6 (6.7, 23.6) µg/g of petroleum by UV/F, values confirmed by the combined $\Sigma 2$, UV/F run. These values were lower than those previously observed in Bay 9 on the 1982 field samples: 25.0 (17.0, 36.0) µg/g, in spite of the higher apparent **levels** of oil seen in the 1983 surface sediments.




FIGURE 3.38. SUMMARY OF OIL CONCENTRATIONS IN Macoma calcaria, by UV/F, ($\mu\text{g/g}$ dry wt.).





FIGURE 3.39. Macoma UV/F SPECTRA.

FIGURE 3.40. REPRESENTATIVE Macoma GC² DETERMINATIONS.




FIGURE 3.41. Macoma AROMATIC PROFILES BY GC²/MS (BAY 11).

3.3.3.2b Oil Composition by GC2. Levels of oil, lower than those observed in Bay 11, were seen in the combined Bay 9 sample. That oil is present is noted by the small quantity of UCM material and the presence of phytane (.02 µg/g) in the sample.

3.3.3.2c Aromatic Hydrocarbon Composition by GC²/MS. Petroleum aromatics were detected in Macoma from Bay 9 in the 1-5 rig/g range with the phenanthrene most abundant. Alkylated phenanthrenes and DBT were detected at levels of up to 150 rig/g in 1982. The absence of these homologous series in the Macoma is unexpected and can not be adequately explained at present.

3.3.3.3 Bay 7

3.3.3.3a Oil Concentrations by UV/F. The Macoma samples from Bay 7 contained detectable quantities of oil in the 1.0 to 5 µg/g range. The combined f2 UV/F analysis yielded an oil concentration value of 4.7 µg/g. The mean of the five tissue samples was 4.4 (3.7, 6.5) µg/g was higher than the 1.9 (1.6, 2.3) µg/g observed in 1982.

3.3.3.3b Oil Composition by GC2. The GC² trace of the combined Bay 7 Macoma (Figure 3.42) sample did reveal the low level presence of UCM and phytane amidst a composition similar to sedimentary hydrocarbons. Biogenic material dominated although the phytane levels of .02 µg/g convert to an equivalent concentration of 2.9 µg/g.

3.3.3.3C Aromatic Hydrocarbon Composition by GC²/MS. Petroleum aromatics in the 2-6 rig/g range were detected in the Bay 7 composite sample. Alkylated naphthalenes and phenanthrenes were the only compounds detected. Bay 7 animals contained 1-20 rig/g in 1982.

3.3.4 Astarte borealis

3.3.4.1 Bay 11

3.3.4.1a Oil Concentrations by UV/F. Concentrations of oil in Astarte (Figures 3.43 and 3.44) from 1983 were 15.2 (6.1, 38.4) µg/g. When last sampled in 1982, these samples averaged 37.0 (33.0, 38.0) µg/g which spanned the 1982 range as well. However, the lower 1983 mean values indicate that oil levels in Astarte are still decreasing.




FIGURE 3.42. Macoma SATURATED HYDROCARBONS FROM BAY 7.







FIGURE 3.43. UV/F SPECTRA OF Astarte FROM BAY 11.

TISSUE PLOTS



^aMean(Lower 95%, Upper 95%)

FIGURE 3.44. SUMMARY OF OIL CONCENTRATIONS IN Astarte borealis, by UV/F, ($\mu\text{g/g}$ dry wt.).

3.3.4.1b Oil Composition by GC2. GC2 analysis of the Astarte composite revealed the same features characteristic of other low level samples: 1) a predominantly biogenic assemblage; 2) UCM material; 3) small quantities of phytane; 4) n-alkanes in the C₂₀ to C₃₀ range. Additionally, those Bay 11 (and other Astarte) contained a double UCM (see Figure 3.45) of unknown origin, but possibly resulting from biodegradation processes.

3.3.4.1c Aromatic Hydrocarbon' Composition by GC²/MS. Low levels of alkylated phenanthrene (1-3 rig/g) and dibenzphtiohenes (~1 rig/g) were detected in the 1983 field samples much lower than the 10-150 rig/g values observed in the 1982 field samples.


3.3.4.2 Bay 9

3.3.4.2a Oil Concentrations by UV/F. Concentrations of oil in Bay 9 Astarte (Figure 3.44) were 7.0 (5.6, 8.7) µg/g. When last sampled in 1982, the corresponding values were 19.0 (10.0, 40.0) µg/g. As in Bay 11, the Astarte values are continuing to decrease despite an input of oil to the sediments of both bays.

3.3.4.2b Oil Composition by GC². The Bay 9 Astarte chromatogram is shown in Figure 3.46. This GC2 trace is enhanced in the vertical direction several times to illustrate several interesting features:

1. The double small unresolved areas of the chromatogram overlaying the larger more frequently encountered UCM.
2. The biogenic olef inic cluster at C₁₈-C₁₉,
3. The n-alkane distribution from C₂₀ to C₃₀ characteristic of petrogenic paraffins,
4. The significant quantity of phytane (0.11 µg/g) equivalent to an 18.3 µg/g concentration of oil residues higher than that detected by UV/F.

3.3.4.2c Aromatic Hydrocarbon Composition by GC²/MS. Alkylated phenanthrenes (C₁-C₃) were detected in the Astarte samples in the 5-10 rig/g range, lower by a factor of four than the levels seen in the previous, 1982, sampling.






FIGURE 3.46. Astarte AROMATIC PROFILES BY GC²/MS (BAY 11).

3.3.4.3 Bay 7

3.3.4.3a Oil Concentrations by UV/F. The Bay 7 Astarte oil concentrations determined by UV/F (Figure 3.44) were 1.7 (.60, 4.8) $\mu\text{g/g}$, less than the values detected in 1982, 6.8 (3.1, 14.8) $\mu\text{g/g}$. This decrease in oil levels parallels the decreases in Bays 11 and 9 for this species.

3.3.4.3b Oil Composition by GC2. Traces of oil are seen in the Bay 7 Astarte GC2 trace (Figure 3.45B). Note the double UCM, as was observed in Bay 9 and the residual n-alkane material in the mid-boiling range ($\text{C}_{20}\text{-C}_{30}$) in the chromatogram. These double UCM features may be sourced during the biodegradation of acquired oil residues in vivo although this is an untested hypothesis. This double UCM may also be an artifact resulting from an unusually complex array of biogenic hydrocarbons in these samples.

3.3.4.3c Aromatic Hydrocarbon Composition by GC²/MS. Individual **alkylated naphthalene** (5-10 rig/g), phenanthrene (25 rig/g) and dibenzothiophene (i O- 15 rig/g) compounds were detected by GC²/MS. No aromatics were previously detected in the 1982 sample set.

3.3.5 Strongylocentrotus droebachiensis (URCHINS)

3.3.5.1 Bay 11

3.3.5.1a Oil Concentrations by UV/F. When last sampled in September, 1982, level of oil in urchins were 67.0 (40.0, 113) $\mu\text{g/g}$. The current 1983 results show that higher levels are clearly present (Figure 3.47) and results are summarized in Figure 3.48. Concentrations are 103 (57.1, 187) $\mu\text{g/g}$, which are similar to levels observed in the spring of 1982.

A very important discrepancy between the UV/F results obtained on the unfractionated extracts and that obtained on the combined f₂ fraction was noted. While the mean oil concentration demonstrated on the unfractionated extracts was 103 $\mu\text{g/g}$, the combined f₂ value was 6.1. The UV/F spectra for the unfractionated animals shows a clearly **visible** "oil peak" at 355 nm. This peak is much decreased in the f₂ fraction after **column** chromatography. (The internal standard is recovered.).




FIGURE 3.47. BAY 11 URCHIN RESULTS: A- UV/F; B- SATURATED HYDROCARBON GC2; C- AROMATIC/UNSATURATED HYDROCARBON GC2.

TISSUE PLOTS

| | | | | |
|-----|-----|------|-----|------|
| 163 | 204 | 67.1 | 110 | 48.3 |
| 1 | 2 | 3 | 4 | 5 |

BAY I I
7m 103(56.8, 187)
6.1 (Combined f 2)

| | | | | |
|------------|-----|-----|------|-----|
| 180 | 130 | 100 | 86.8 | 376 |
| 1 | 2 | 3 | 4 | 5 |

BAY 9
7m 150(83.9, 269)
6.8 (Combined f2)

| | | | | |
|-------------|------------|-------------|-------------|------------|
| 18.7 | 5.6 | 28.0 | 11.8 | 7.7 |
| 1 | 2 | 3 | 4 | 5 |

BAY 7
7m 12.2(6.4, 23.2)
.86 (Combined f₂)

^aMean(Lower 95%, Upper 95%)

FIGURE 3.48. SUMMARY OF OIL CONCENTRATIONS IN *Strongylocentrotus droebachiensis*, by UV/F, ($\mu\text{g/g}$ dry wt.).

3.3.5. 1b Oil Composition by GC2. Weathered oil residues are apparent in the GC2 trace shown in Figure 3.47B, where petrogenic phytane, pristane and UCM are readily apparent. No prominent *n*-alkane distributions are observed in the C₂₀ to C₃₀ range for this species.

Aromatic hydrocarbon GC2 traces (e.g. Figure 3.47C) indicate the low relative quantity of petroleum aromatics compared to biogenic olefins.

3.3.5. 1c Aromatic Hydrocarbons by GC²/MS. Large quantities of alkylated phenanthrenes were detected in the Bay 1 urchins. The values 10-190 rig/g are roughly equal to those detected on the 1982 samples. The composition, however, (Figure 3.49) is notably lacking in dibenzothiophene compounds.

3.3.5.2 Bay 9

3.3.5.1a Oil Concentrations by UV/F. UV/F determined oil concentrations on total extracts from Bay 9 were 150 (83.9, 269) µg/g. In spite of the apparent unambiguous quantification of these extracts (Fig 3.50) the UV/F value of the combined f2 fraction yielded a value of 6.8 µg/g. This is similar to the Bay 11 discrepancy. Note, however, that such comparisons were not made in previous years.

The comparable 1982 field sample result 46.0 (25.0, 86.0) µg/g is less than the 1983 values determined on the total extracts. Note that values as high as 760 µg/g were found in May, 1982 in Bay 9 urchins.

3.3.5.2b Oil Composition by GC2. The GC2 trace of the Bay 9 sample contained mainly biogenic compounds although a significant amount of UCM material was detected. The profile was similar in composition to that from Bay 11 (Figure 3.46) although of lesser concentration.

3.3.5.2c Aromatic Hydrocarbon Composition by GC²/MS. The GC²/MS analysis of the Bay 9 urchin's aromatic fraction failed to detect (>1 rig/g) any petroleum aromatics although significant quantities (15-150 rig/g) of four and five ringed aromatics were detected.

3.3.5.3 Bay 7

3.3.5.3a Oil Concentrations by UV/F. Although a significant discrepancy still exists between the UV/F total extract values, 12.2 (6.4, 23.2) µg/g and the UV/F of the




FIGURE 3.49. Strongylocentrotus AROMATIC PROFILES BY GC²/MS (BAY 11).




FIGURE 3.50. Strongylocentrotus UV/F SPECTRA, BAY

fractionated extract (0.9 $\mu\text{g/g}$), the Bay 7 values are much lower than those from Bays 9 and 11. Last year's values were 4.6 (2.3, 9.2) $\mu\text{g/g}$.

3.3.5.3b Oil Composition by GC2. Biogenic compounds dominated the saturated hydrocarbon GC2 determination (Fig. 3.51) although petrogenic inputs are indicated by the presence of a UCM distribution and by small quantities of phytane.

3.3.5.3c Aromatic Hydrocarbon Composition by GC²/MS. Small quantities 1-5 ng/g of C₀ and C₁ phenanthrene were detected in the composite sample. This finding is consistent with the GC2 saturated composition which indicates that sediment material was present in the urchin tissues. C₀P and C₁P are common low level background components of sediments.

3.3.6 MILNE INLET ANIMALS

One composite sample of each species was obtained from the benthos of Milne Inlet and the western side (non-spill side) of Ragged Island. These animals were analyzed by UV/F, GC² and GC²/MS.

UV/F results on the total extracts are as follows:

| | | |
|----------------|---|----------------------|
| <u>Mya</u> | = | <0.8 $\mu\text{g/g}$ |
| <u>Serpis</u> | = | <0.8 |
| <u>Macoma</u> | = | <0.8 |
| <u>Astarte</u> | = | 1.3 |
| Str. | = | <0.8 |

GC² results detected only biogenic components, mainly pristane and no UCM material. Therefore, no petrogenic components were detected.

GC²/MS results were as follows:

| | | |
|----------------|---|---|
| <u>Mya</u> | : | Phenanthrene detected at 1 ng/g |
| <u>Serpis</u> | : | none detected |
| <u>Macoma</u> | : | naphthalene = 2 ng/g phenanthrene = 4 ng/g |
| <u>Astarte</u> | : | naphthalene = 1 ng/g phenanthrene = 1 ng/g |
| Str. | : | phenanthrene = 1 ng/g |




FIGURE 3.51. GC2 TRACE OF Strongylocentrotus BAY 7 SATURATED HYDROCARBONS

3.4 Summary of Temporal Trends in Mean Oil Concentrations in Sediment and Animals

Details of the distributions of petroleum hydrocarbons in sediments and **benthic animals** determined from the 1983 sampling have been presented in Sections 3.2 and 3.3. To put these findings in temporal perspective, the 1980-1983 trends in petroleum residues for sediments and animals of Bay 11 are presented in Figures 3.52 and 3.53. These are geometric mean concentrations obtained from **log-transf** ormed concentration data. Confidence limits are defined in Sections 3.2 and 3.3. The important features of these plots are:

1. The general increase in sediment levels at all stations. Note that the 3m **benthic** transect (No. 3) value is off scale in Figure 3.52.
2. The August 1982 to August 1983 increase (or leveling off) of oil concentrations in Strongylocentrotus and Macoma. Note that the initial increase in oil levels (September 1981) for all animals is believed to have been caused by intrusion of dispersed oil from the Bay 9 release, not from the Bay 11 surface release.

The Bay 9 results are summarized in Figures 3.54 and 3.55. The important features to note are:

1. After an initial increase in Bay 9 sediment oil levels, values decreased in 1982 followed by a general increase, except at Benthic Transect No. 1 at 3m, in 1983. Note that oil levels remained elevated (approximately 4-6 ppm) in the 10m deep microbiology plots in 1982.
2. A long-term deputation is seen to occur in most of the Bay 9 **benthic** animals. However, after two years, oil levels in Macoma and Astarte are still 15-20 times background levels, while Strongylocentrotus has apparently acquired significantly more oil between 1982 and 1983 paralleling the increase in sediment levels.




FIGURE 3.52. BAY 11 SEDIMENT OIL CONCENTRATIONS (1980-1983) ($\mu\text{g/g}$; BY UV/F)




FIGURE 3.53. OIL CONCENTRATIONS IN ANIMALS: BAY 11 (μg/g; BY UV/F) (1980-1983).




FIGURE 3.54. BAY 9 SEDIMENT OIL CONCENTRATIONS (1980-1983) ($\mu\text{g/g}$; BY UV/F)




FIGURE 3.55. OIL CONCENTRATIONS IN ANIMALS: BAY 9 (µg/g; BY UV/F) 1980-1983).

Additional temporal information is presented for Bay 7 benthic animals (Figure 3.56). Note that the sediment levels in Bay 7 while still uniformly low in 1983 did apparently increase to about 2 ppm on the average in 1983. These are very low, but detectable levels which may be affecting Strongylocentrotus (Figure 3.56) in Bay 7 in 1983.

Bay 10 benthic animal temporal trends (Figure 3.57) were only followed through the 1982 field season.




FIGURE 3.56. OIL CONCENTRATIONS IN BENTHIC ANIMALS: BAY 7 (µg/g; BY UV/F) (980-1983).




FIGURE 3.57. OIL CONCENTRATIONS IN ANIMALS: BAY 10 ($\mu\text{g/g}$; BY UV/F) (1980-1982)

SECTION FOUR

SHORELINE STUDY

A detailed study of the oil concentrations and compositions in a set of 96 beach sediment samples was conducted. A summary of the samples analyzed is shown in Table 4.1.

All samples were extracted and the extracts fractionated by column chromatography to yield a saturated and aromatic hydrocarbon fraction for each sample. Concentrations of petroleum hydrocarbons were determined by **microgravimetry** of these fractions. The total extractable material was also weighed. This latter value represents the total petroleum value which includes the hydrocarbons **plus** polar material (non-hydrocarbon) in the stranded petroleum.

The saturated hydrocarbon composition was determined on **all samples by GC²** analysis. Two ratios best describe the **GC²-determined** composition:

1. The SHWR reflects the weathering status due mainly to physical-chemical processes (i.e. evaporation, dissolution). As the n-C **10** to n-C **17** hydrocarbons are lost due to those processes, the SHWR approaches zero from an unweathered value of **~2.5**. Note that where petroleum (kerosene) based dispersants have been used as part of the cleanup experiment, **alkane** components added in the C **10** to C **17** (from kerosene) range create SHWR > 2.5. Where SHWR exceeds that in the crude oil itself, kerosene additions are the likely explanation.
2. **ALK/ISO** ratio - In the Lagomedio oil, the ratio of **n-alkanes** to **isoprenoid alkanes** in the C **13** - C **19** boiling region is **~2.5**. Normal **alkanes** are the most readily biodegraded components. Thus, as biodegradation proceeds, this ratio approaches zero, and the GC² trace becomes progressively depleted in **n-alkanes**. This is not to say that the **isoprenoids** themselves are not degraded; however, they are degraded more slowly than are the **n-alkanes**. Additions of kerosene or **biogenic** materials can confound the interpretation of this ratio. Kerosene (and diesel) additions add components in the boiling range of the components which make up this ratio. Also, at low levels of oil the inputs of **biogenic pristane** (one of the isoprenoids) and n-C **15** and n-C **17** (from plankton; two of the **n-alkanes**) can affect this ratio.

TABLE 4.1. SHORELINE STUDY HYDROCARBON CHEMISTRY RESULTS SAMPLE DESCRIPTIONS

| Test Year | Sample ID | Plot | Depth | Date |
|----------------|-----------|-------------------|------------------------|----------|
| 1980 | 4050 | Bay 102 Oil Patch | | 83-08-17 |
| 1980 | 4051 | L1 | Upper Surface | 83-08-17 |
| 1980 | 4052 | L1 | Upper Sub-Surface | 83-08-17 |
| 1980 | 4053 | L1 | Lower Surface | 83-08-17 |
| 1980 | 4054 | L1 | Lower Sub-Surface | 83-08-17 |
| 1980 | 4055 | L2 | Surface | 83-08-17 |
| 1980 | 4056 | L2 | Sub-Surface | 83-08-17 |
| 1980 | 4057 | T1 | Surface | 83-08-17 |
| 1980 | 4058 | T1 | Sub-Surface | 83-08-17 |
| 1980 | 4059 | T2 | Surface | 83-08-17 |
| 1980 | 4060 | T2 | Sub-Surface | 83-08-17 |
| 1980 | 4957 | T2 | Upper Sub-Surface | 83-08-17 |
| 1981 | 4120 | CC | Surface | 83-08-17 |
| 1981 | 4121 | CC | Sub-Surface | 83-08-17 |
| 1981 | 4122 | CE | Surface | 83-08-17 |
| 1981 | 4123 | CE | Sub-Surface | 83-08-17 |
| Ragged Channel | 4124 | Bay 9 | 100, Upper Surface | 83-08-10 |
| Beaches | 4126 | Bay 9 | 100, Mid Surface | 83-08-10 |
| Beaches | 4127 | Bay 9 | 100, Mid-Sub-Surface | 83-08-10 |
| Beaches | 4128 | Bay 9 | 100, Lower Surface | 83-08-10 |
| Beaches | 4129 | Bay 9 | 100, Lower Sub-Surface | 83-08-10 |
| Beaches | 4130 | Bay 9 | 300, Mid-Surface | 83-08-10 |
| Beaches | 4132 | Bay 9 | 300, Lower Surface | 83-08-10 |
| Beaches | 4134 | Bay 11 | 2, Upper Surface | 83-08-16 |
| Beaches | 4135 | Bay 11 | 6, Upper Surface | 83-08-16 |
| Beaches | 4136 | Bay 11 | 2, Mid-Surface | 83-08-16 |
| Beaches | 4137 | Bay 11 | 6, Mid-Surface | 83-08-16 |
| Beaches | 4138 | Bay 11 | 2, Lower Surface | 83-08-16 |
| Beaches | 4139 | Bay 11 | 6, Lower Surface | 83-08-16 |
| Beaches | 4140 | Bay 11 | 4, Upper Surface | 83-08-16 |
| Beaches | 4141 | Bay 11 | 8, Upper Surface | 83-08-16 |
| Beaches | 4142 | Bay 11 | 4, Mid-Surface | 83-08-16 |
| Beaches | 4143 | Bay 11 | 8, Mid-Surface | 83-08-16 |
| Beaches | 4144 | Bay 11 | 4, Lower Surface | 83-08-16 |
| Beaches | 4145 | Bay 11 | 8, Lower Surface | 83-08-16 |
| Beaches | 4146 | Bay 11 | X1 Surface | 83-08-16 |
| Beaches | 4147 | Bay 11 | X2 Surface | 83-08-16 |
| Beaches | 4148 | Bay 11 | X3 Surface | 83-08-16 |
| Beaches | 4149 | Bay 11 | X4 Surface | 83-08-16 |
| Beaches | 4150 | Bay 11 | X5 Surface | 83-08-16 |

TABLE 4.1.
(Continued)

| Test Year | Sample ID | Plot | Depth | Date |
|----------------|-----------|-----------------|--------------------|----------|
| Ragged Channel | 4946 | Bay 11 | X 1 Sub-Surface | 83-08-16 |
| Beaches | 4947 | Bay 11 | X2 Sub-Surface | 83-08-16 |
| Beaches | 4948 | Bay 11 | X3 Sub-Surface | 83-08-16 |
| Beaches | 4949 | Bay 11 | X4 Sub-Surface | 83-08-16 |
| Beaches | 4950 | Bay 11 | X5 Sub-Surface | 83-08-16 |
| Beaches | 4951 | Bay 11 | X6 Sub-Surface | 83-08-16 |
| Beaches | 4952 | Bay 9 | 300, Upper Surface | 83-08-16 |
| Beaches | 4955 | Crude Oil Point | X7 Surface | 83-08-17 |
| Beaches | 4956 | Crude Oil Point | X8 Sub-Surface | 83-08-17 |
| 1982 | 4301 | ICC | Surface | 83-08-17 |
| 1982 | 4302 | ICC | Sub-Surface | 83-08-17 |
| 1982 | 4303 | ICE | Surface | 83-08-17 |
| 1982 | 4304 | ICE | Sub-Surface | 83-08-17 |
| 1982 | 4305 | IDEC | Surface | 83-08-17 |
| 1982 | 4306 | IDEC | Sub-Surface | 83-08-17 |
| 1982 | 4307 | IDEE | Surface | 83-08-17 |
| 1982 | 4308 | IDEE | Sub-Surface | 83-08-17 |
| 1982 | 4309 | IDBC | Surface | 83-08-17 |
| 1982 | 4310 | IDBC | Sub-Surface | 83-08-17 |
| 1982 | 4311 | IDBE | Surface | 83-08-17 |
| 1982 | 4312 | IDBE | Sub-Surface | 83-08-17 |
| 1982 | 4313 | IMC-C | Berm Surface | 83-08-17 |
| 1982 | 4314 | IMC-C | Berm Sub-Surface | 83-08-17 |
| 1982 | 4315 | IMC-M | Berm Surface | 83-08-17 |
| 1982 | 4316 | IMC-M | Berm Sub-Surface | 83-08-17 |
| 1982 | 4317 | IMC-C | Back Surface | 83-08-17 |
| 1982 | 4318 | IMC-C | Back Sub-Surface | 83-08-17 |
| 1982 | 4319 | IMC-M | Back Surface | 83-08-17 |
| 1982 | 4320 | IMC-M | Back Sub-Surface | 83-08-17 |
| 1982 | 4321 | IME-C | Berm Surface | 83-08-17 |
| 1982 | 4322 | IME-C | Berm Sub-Surface | 83-08-17 |
| 1982 | 4323 | IME-M | Berm Surface | 83-08-17 |
| 1982 | 4324 | IME-M | Berm Sub-Surface | 83-08-17 |
| 1982 | 4325 | IME-C | Back Surface | 83-08-17 |
| 1982 | 4326 | IME-C | Back Sub-Surface | 83-08-17 |
| 1982 | 4327 | IME-M | Back Surface | 83-08-17 |
| 1982 | 4328 | IME-M | Back Sub-Surface | 83-08-17 |
| Norwegian | 4329 | N1, T2 West | Surface | 83-08-17 |
| Norwegian | 4330 | N2, T2 West | Composite | 83-08-17 |
| Norwegian | 4331 | N3, T2 East | Surface | 83-08-17 |
| Norwegian | 4332 | N4, T2 East | Composite | 83-08-17 |
| Norwegian | 4333 | N5, 102E | | 83-08-17 |
| Norwegian | 4334 | N6, 102F | | 83-08-17 |
| Norwegian | 4335 | N7, 102G | | 83-08-17 |
| Norwegian | 4336 | N8, 102H | | 83-08-17 |

TABLE 4.1.
(Continued)

| Test Year | Sample ID | Plot | Depth | Date |
|--------------|--------------|-----------|-----------|----------|
| Norwegian | 4337 | N9, 102A | | 83-08-17 |
| Norwegian | 4338 | N10, 102B | | 83-08-17 |
| Norwegian | 4339 | Nil, 106A | Surface | 83-08-17 |
| Norwegian | 4340 | N12, 106A | Composite | 83-08-17 |
| Norwegian | 4341 | N13, 106C | Surface | 83-08-17 |
| Norwegian | 4342 | N14, 106C | Composite | 83-08-17 |
| Norwegian | 4343 | N15, 106D | Surface | 83-08-17 |
| Norwegian | 4344 | N16, 106D | Composite | 83-08-17 |
| Norwegian | 4345 | N17, 106E | Surface | 83-08-17 |
| Norwegian | 4346 | N18, 106E | Composite | 83-08-17 |

Several examples illustrate these points. The GC2 traces in Figure 4. illustrate:

1. Figure 4.1a, a relatively unweathered oil sample; SHWR = 2.0; ALK/ISO = 2.1; AWR (see below) = 3.0 (S4052; L 1, upper sub-surface; 8960 ppm).
2. Figure 4.1b - A moderately weathered oil sample; physical-chemical weathering due to evaporation, - SHWR = 1.1; little biodegradation, - ALK/ISO = 1.9 (S4057; T1; surface; 33800 ppm). Abundant UCM present.
3. Figure 4.1c - A sample to which a low boiling distillate has been added; SHWR = 12.0; ALK/ISO = 3.7 (S4309); IDBC; surface; 1620 ppm).

The GC2 traces in Figure 4.2 indicate:

1. Figure 4.2a - A moderately weathered (SHWR = 2.0) biodegraded (ALK/ISO = 0.7) oil sample (S4121; CC; sub-surface; 300 ppm). UCM present, bimodal.
2. Figure 4.2b - A sample which has received significant quantities of light distillate (e.g., kerosene or diesel); SHWR = 3.3, ALK/ISO = 3.2; (S4120 ; CC; surface; 130 ppm).

The results shown in Figure 4.3 illustrate:

1. Figure 4.3a - A relatively unweathered, unbiodegraded sample; SHWR = 1.8; ALK/ISO = 2.2 (S4330; N2, T2 west, composite, 28,600 ppm).
2. Figure 4.3b - A weathered (SHWR = 1.1) but unbiodegraded sample (ALK/ISO = 2.1) (S4332; N4, T2, east, composite; 36,300 ppm).
3. Figure 4.3c - A moderately weathered (SHWR = 1.5) biodegraded (ALK/ISO = 0.9) sample (S4340; N 12, 106C, surface, 39,700 ppm). Bimodal UCM present.

Illustrations of biodegraded, weathered oil, the Bay 11 beach, have previously been shown in Section 3.2.1.7. (Figure 3.21).

The quantitative and compositional results on the shoreline sediments are summarized in Tables 4.2 through 4.6 for the various sample sets. These data include the gravimetric (oil concentrations) and GC2 (saturated hydrocarbon compositions) as well as GC²/MS determinations of the aromatic weathering ratio (AWR) in a selected set of




FIGURE 4.1. REPRESENTATIVE SHORELINE SEDIMENT SATURATED HYDROCARBON GC² DETERMINATIONS A- S4052, LI; B- 4057, TI; C- 4309, IDBC.




FIGURE 4.2. REPRESENTATIVE SHORELINE SEDIMENT SATURATED HYDROCARBON GC² DETERMINATIONS A- 4124, CC; 54120, CC.



REPRESENTATIVE SHORELINE SEDIMENT SATURATED HYDROCARBON GC² DETERMINATIONS: A-S4330; B-S4332; C-S4340.

TABLE 4.2. SHORELINE STUDY 1983 HYDROCARBON CHEMISTRY RESULTS - ANALYTICAL **RESULTS**; 1980 TEST PLOTS.

| Plot | Depth | Saturated Hydrocarbon µg/g | Aromatic Hydrocarbon µg/g | Total Hydrocarbon µg/g | Total Extractable µg/g | SHWR | ALK/ISO | AWR |
|-----------------------|-------------------|----------------------------------|---------------------------------|------------------------------|------------------------------|------|---------|-----|
| Bay 102, Oil Patch | | 889. | 526. | 1,940. | 1,930. | 1.2 | 0.8 | |
| L1 | Upper Surface | 686. | 345. | 1,030. | 1,470. | 1.2 | 0.9 | 1.3 |
| L1 | Upper Sub-Surface | 4,890. | 2,460. | 7,350. | 8,960. | 2.0 | 2.1 | 3.0 |
| L1 | Lower Surface | 18.3 | 9.4 | 27.7 | 55.0 | 1.0 | 1.7 | 1.5 |
| L1 | Lower Sub-Surface | 0.0 | 1.9 | 1.9 | 16.0 | 1.2 | 1.0 | |
| L2 | Surface | 4.2 | 5.1 | 9.3 | 34.2 | 1.3 | 1.7 | |
| L2 | Sub-Surface | 2.0 | 0.9 | 2.9 | 23.7 | 1.4 | 1.1 | |
| T1 | Surface | 2,780. | 1,430. | 4,210. | 6,450. | 1.1 | 1.9 | |
| T1 | Sub-Surface | 6,170. | 4,250. | 10,400. | 13,900. | 1.8 | 2.2 | |
| T2 | Surface | 17,100. | 10,500. | 27,600. | 33,800. | 1.4 | 2.2 | |
| T2 | Sub-Surface | 9,590. | 5,470. | 15,100. | 19,900. | 2.2 | 2.2 | |
| T2 | Upper Sub-Surface | 242. | 159. | 401. | 635. | 1.2 | 0.2 | |

TABLE 4.3. SHORELINE STUDY 1983 **HYDROCARBON CHEMISTRY RESULTS - ANALYTICAL RESULTS**; 1981 TEST PLOTS.

| Plot | Depth | Saturated Hydrocarbon µg/g | Aromatic Hydrocarbon µg/g | Total Hydrocarbon µg/g | Total Extractable µg/g | SHWR | ALK/ISO | AWR |
|------|-------------|----------------------------------|---------------------------------|------------------------------|------------------------------|------|---------|-----|
| CC | Surface | 33.7 | 19.8 | 53.5 | 131. | 3.3 | 3.2 | |
| c c | Sub-Surface | 117. | 78.6 | 196. | 299. | 2.0 | 0.7 | |
| CE | Surface | 6.9 | 5.6 | 12.5 | 62.0 | 2.0 | 2.1 | |
| CE | Sub-Surface | 18.1 | 14.7 | 32.8 | 78.5 | 1.9 | 1.4 | |

TABLE 4.4. SHORELINE STUDY 1983 HYDROCARBON CHEMISTRY RESULTS - ANALYTICAL RESULTS; 1982 TEST PLOTS.

| Plot | Depth | Saturated Hydrocarbon µg/g | Aromatic Hydrocarbon µg/g | Total Hydrocarbon µg/g | Total Extractable µg/g | SHWR | ALK/ISO | AWR |
|-------|------------------|----------------------------------|---------------------------------|------------------------------|------------------------------|------|---------|-----|
| ICC | Surface | 112. | 73.6 | 185. | 437. | 1.6 | 1.4 | 1.6 |
| ICC | Sub-Surf ace | 2.5 | 1.8 | 4.3 | 55.3 | 1.5 | 1.9 | |
| ICE | Surface | 356. | 214. | 570. | 924. | 1.5 | 1.7 | 1.5 |
| ICE | Sub-Surface | 1.8 | 4.7 | 6.5 | 230. | 1.8 | 0.6 | |
| IDEC | Surface | 276. | 163. | 439. | 709. | 1.4 | 2.0 | |
| IDEC | Sub-Surf ace | 1.4 | 2.6 | 4.0 | 75.5 | 1.1 | 2.5 | |
| IDEE | Surface | 297. | 188. | 484. | 887. | 1.4 | 1.5 | |
| IDEE | Sub-Surface | 3.5 | 9.5 | 13.0 | 332. | 2.5 | 0.4 | |
| IDBC | Surface | 523. | 102. | 624. | 1,620. | 12. | 3.7 | |
| IDBC | Sub-Surface | 380. | 54.7 | 435. | 668. | 17. | 3.6 | |
| IDBE | Surface | 354. | 75.6 | 429. | 868. | 14. | 3.9 | |
| IDBE | Sub-Surf ace | 2.7 | 6.2 | 8.9 | 121. | 2.8 | 1*9 | |
| IMC-C | Berm. Surface | 19,600. | 11,600. | 31,200. | 45,200. | 1.5 | 2.2 | 2.3 |
| IMC-C | Berm Sub-Surface | 3,860. | 2,800. | 6,660. | 9,480. | 2.4 | 2.3 | |
| IMC-M | Berm Surf ace | 19,300. | 12,000. | 31,300. | 41,200. | 1.6 | 2.3 | |
| IMC-M | Berm Sub-Surface | 5,960. | 3,490. | 9,460. | 11,800. | 2.1 | 2.3 | |
| IMC-C | Back Surface | 17,100. | 10,500. | 27,600. | 41,000. | 1.4 | 1.9 | 1.7 |
| IMC-C | Back Sub-Surface | 118. | 87.8 | 206. | 376. | 1.9 | 2.1 | 3.1 |
| IMC-M | Back Surface | 8,230. | 5,680. | 13,900. | 15,700. | 1.7 | 2.2 | |
| IMC-M | Back Sub-Surface | 929. | 679. | 1,610. | 2,410. | 2.0 | 2.1 | |
| IME-C | Berm Surface | 6210. | 3,790. | 10,000. | 11,900. | 1.7 | 2.1 | 2.8 |
| IME-C | Berm Sub-Surface | 5,170. | 2,860. | 8,020. | 9,670. | 2.0 | 2.3 | |
| IME-M | Berm Surface | 6,070. | 3,520. | 9,600. | 11,900. | 1.7 | 2.3 | |
| IME-M | Berm Sub-Surface | 6,440. | 3,650. | 10,100. | 11,700. | 2.0 | 2.2 | |
| IME-C | Back Surface | 20,900. | 13,300. | 34,200. | 44,100. | 1.5 | 1.4 | 1.8 |
| IME-C | Back Sub-Surface | 389. | 259. | 648. | 947. | 1.6 | 2.0 | 2.0 |
| IME-M | Back Surface | 13,400. | 6,000. | 19,400. | 20,400. | 1.6 | 2.4 | |
| IME-M | Back Sub-Surface | 4,890. | 3,300. | 8,190. | 12,900. | 2.1 | 2.3 | |

12E

TABLE 4.5. SHORELINE STUDY 1983 HYDROCARBON CHEMISTRY RESULTS - ANALYTICAL RESULTS; RAGGED CHANNEL BEACHES.

| Plot | Depth | Saturated Hydrocarbon µg/g | Aromatic Hydrocarbon µg/g | Total Hydrocarbon µg/g | Total Extractable µg/g | SHWR | ALK/ISO | AWR |
|------------------------|-----------------------------|----------------------------------|---------------------------------|------------------------------|------------------------------|------------|------------|------------|
| Bay 9 | 100, Upper Surface | 0.0 | 1.0 | 1.0 | 8.1 | 2.0 | 0.8 | 1.2 |
| Bay 9 | 100, Mid-Surface | 0.9 | 0.4 | 1.3 | 6.0 | 1.2 | 1.4 | |
| Bay 9 | 100, Mid Sub-Surface | 0.0 | 0.0 | 0.0 | 11.3 | 1.7 | 0.9 | 1.6 |
| Bay 9 | 100, Lower Surface | 0.2 | 0.0 | 0.2 | 2.92 | 0 | 0 | |
| Bay 9 | 100, Lower Sub-Surface | 0.8 | 0.8 | 1.6 | 11.6 | 1.0 | 0.9 | |
| Bay 9 | 300, Mid-Surface | 0.5 | 1.1 | 1.6 | | 1.2 | 0.8 | |
| Bay 9 | 300, Lower Surface | 0.0 | 0.7 | 0.7 | 9.7 | 1.2 | 1.6 | |
| Bay 11 | 2, Upper Surface | 221. | 180. | 401. | 1,260. | 1.0 | 1.0 | 1.0 |
| Bay 11 | 6, Upper Surface | 8,490. | 5,380. | 13,900. | 17,300. | 1.6 | 1.9 | 2.2 |
| Bay 11 | 2, Mid-Surface | 601. | 307. | 908. | 1,880. | 1.1 | 0.4 | 1.1 |
| Bay 11 | 6, Mid-Surface | 9,210. | 4,280. | 13,500. | 16,700. | 1.7 | 2.1 | |
| Bay 11 | 2, Lower Surface | 55.1 | 16.9 | 72.0 | 197. | 1.1 | 0.2 | |
| Bay 11 | 6, Lower Surface | 2,580. | 1,310. | 3,890. | 6,160. | 1.0 | 0.3 | |
| Bay 11 | 4, Upper Surface | 12,200. | 7,230. | 19,400. | 25,400. | 1.6 | 2.5 | |
| Bay 11 | 8, Upper Surface | 2,380. | 1,100. | 3,480. | 4,810. | 1.1 | 1.4 | |
| Bay 11 | 4, Mid-Surface | 4,220. | 1,930. | 6,150. | 10,800. | 1.0 | 0.7 | |
| Bay 11 | 8, Mid-Surface | 1,010. | 503. | 1,510. | 2,450. | 1.0 | 1.4 | |
| Bay 11 | 4, Lower Surface | 29.6 | 1.1 | 42.7 | 94.0 | 1.2 | 0.4 | |
| Bay 11 | 8, Lower Surface | 835. | 422. | 1,260. | 2,100. | 1.1 | 0.3 | |
| Bay 11 | X 1 Surface | 16,200. | 1,050. | 17,200. | 26,900. | 2.0 | 2.2 | 2.5 |
| Bay 11 | X2 Surface | 216. | 107. | 322. | 600. | 1.3 | 1.2 | |
| Bay 11 | X3 Surface | 332. | 186. | 518. | 809. | 1.8 | 2.1 | |
| Bay 11 | X4 Surface | 122. | 69.8 | 192. | 730. | 1.6 | 1.8 | 2.1 |
| Bay 11 | X 5 Surface | 7,670. | 5,130. | 12,800. | 15,900. | 1.4 | 1.0 | |
| Bay 11 | X 1 Sub-Surface | 4,640. | 2,380. | 7,020. | 8,980. | 1.6 | 2.0 | 2.3 |
| Bay 11 | X2 Sub-Surface | 41.4 | 18.6 | 78.6 | 151.0 | 1.2 | 0.7 | |
| Bay 11 | X3 Sub-Surface | 2.4 | 2.3 | 4.7 | 72.2 | 1.3 | 1.4 | |
| Bay 11 | X4 Sub-Surface | 0.9 | 3.0 | 3.9 | 227. | 1.3 | 1.4 | 3.5 |
| Bay 11 | X5 Sub-Surface | 3,430. | 2,180. | 5,600. | 7,180. | 1.8 | 1.8 | |
| Bay 11 | X6 Sub-Surface | 164. | 105. | 269. | 688. | 1.7 | 2.1 | |
| Bay 11 | 300, Upper Surface | 0.7 | 0.2 | 0.9 | 10.8 | 1.1 | 1.5 | |
| Crude Oil Point | X7 Surface | 1,060. | 467. | 1,530. | 2,600. | 1.0 | 1.2 | |
| Crude Oil Point | X8 Sub-Surface | 1,220. | 724. | 1,940. | 3,000. | 1.0 | 1.6 | |

TABLE 4.6. SHORELINE STUDY 1983 HYDROCARBON CHEMISTRY RESULTS - ANALYTICAL RESULTS; NORWEGIAN TEST PLOTS.

| Plot | Depth | Saturated Hydrocarbon µg/g | Aromatic Hydrocarbon µg/g | Total Hydrocarbon µg/g | Total Extractable µg/g | SHWR | ALK/ISO | AWR |
|--------------|-----------|----------------------------------|---------------------------------|------------------------------|------------------------------|------|---------|-----|
| N 1, T2 West | Surface | 34,200. | 14,200. | 48,400. | 72,600. | 1.4 | 2.0 | |
| N2, T2 West | Composite | 16,300. | 9,140. | 25,400. | 28,600. | 1.8 | 2*2 | |
| N3, T2 East | Surface | 18,200. | 9,830. | 28,100. | 35,400. | 1.3 | 2.1 | |
| N4, T2 East | Composite | 18,300. | 9,950. | 28,300. | 36,300. | 1.1 | 2.1 | |
| N5, 102 E | | 5,490. | 3,250. | 8,740. | 11,300. | 1.4 | 1.2 | |
| N6, 102 F | | 5,470. | 3,690. | 9,160. | 11,700. | 1.8 | 2.1 | |
| N7, 102 G | | 10,700. | 7,210. | 17,900. | 21,100. | 1.8 | 2.2 | |
| N8, 102 H | | 18,700. | 11,800. | 30,500. | 35,800. | 1.7 | 1.8 | |
| N9, 102 A | | 12,700. | 6,870. | 19,500. | 19,000. | 1.6 | 2.1 | |
| N10, 102 B | | 11,900. | 5,920. | 17,900. | 20,900. | 1.5 | 2.1 | |
| N11, 106 A | Surface | 29,200. | 18,100. | 47,300. | 66,100. | 1.3 | 1.1 | |
| N12, 106 A | Composite | 5,180. | 3,700. | 8,880. | 11,600. | 1.5 | 0.9 | |
| N13, 106 C | Surface | 24,400. | 12,500. | 36,800. | 39,700. | 1.4 | 2.0 | |
| N14, 106 C | Composite | 4,370. | 2,380. | 6,750. | 8,970. | 1.5 | 2.0 | |
| N15, 106 D | Surface | 25,000. | 13,900. | 38,800. | 45,400. | 1.4 | 1.5 | |
| N16, 106 D | Composite | 13,400. | 7,770. | 21,100. | 24,900. | 1.7 | 1.6 | |
| N17, 106 E | Surface | 9,660. | 5,870. | 15,500. | 20,000. | 1.4 | 2.0 | |
| N18, 106 E | Composite | 12,500. | 7,360. | 19,800. | 20,600. | 1.8 | 2.2 | |

samples. The AWR in the spilled oil was 3.5. As light aromatics are removed due to physical chemical weathering, this ratio decreases. Twenty such aromatic fractions were analyzed by GC²/MS.

All of these results must be viewed as the latest in a time series of quantitative and qualitative results on the various test plots.

SECTION 5

DISCUSSION OF RESULTS

Two years after the Ragged Channel (nearshore) spills, the focus of the analytical **biogeochemical** studies is primarily on the movement of oil sourced from the stranded oil on the Bay 11 beach and its impact on the **benthic biota**.

In the summer of 1982, the **oil levels** in the **subtidal** sediments of Bay 11 had increased to 3.0 $\mu\text{g/g}$ (range 0.7-66) at 3 meters and 5.3 $\mu\text{g/g}$ (range 1.3-50) at 7 meters. One of the major findings of the 1983 study was the marked increase in **subtidal** sediment **oil levels** in Bay 11. Oil levels in sediments increased by a factor of 5-10 due to erosion of beached oil and deposition offshore to at least the 7m water depth contour. The microbiology transect in Bay 11 confirms ~~that the bulk of the deposited offshore oil lies in the 3m to 7m depth range and the highest concentrations are found at the south end of the 3m line.~~ However, we have confirmed that 1-8 ppn of oil have been transported further off shore in the deeper (35m) areas in the Bay 11/12 region.

Bay 9 sediment petroleum hydrocarbon values increased between 1982 and 1983. The source of elevated quantities of sedimented oil in the sediments of Bay 9 probably lies in the sediments just south of the Bay 9 study area. Large quantities of oil were observed to be transported from the **oil/dispersant** diffuser system south along the shoreline during the discharge in Bay 9 in 1981. It is possible that elevated oil levels in sediments resulting from this southerly transport may have eventually resulted in infiltration of these sediments into the study area of Bay 9, just north of the diffuser site. Values of 0.6-7 ppm of oil have been detected at 3m in Bay 9 and 6-10 ppm detected at 7 meters. On the average, this represents a threefold increase at 7m and at 3m in Bay 9.

Levels of oil in the Bay 7 sediments have increased slightly over the low levels (1-2 ppm) observed previously. One "hot spot" at 13 ppm was found which also corresponded to higher oil levels in one benthic animal species.

Judging from the compositional profiles of the **subtidal** sedimented oil in Bay 11, the residues are more highly weathered than they had been in 1982. The oil eroded off of the Bay 11 beach is **compositionally** very variable. It appears that weathering (i.e. the combined processes of evaporation, dissolution and biodegradation) is proceeding more rapidly in the lower end of the beach transects. Relatively unweathered oil is still present

in concentrated patches on the Bay 11 beach. However, for the most part it appears that both the oil leached into the water column (i.e. the slick samples) and the sedimented oil (floe samples) are largely depleted in low boiling alkanes (n-C 10 to n-C 15) and aromatics (alkylated benzenes and naphthalenes).

The surface floe hydrocarbon levels in the Bay 3 meter samples were almost twice as high, on the average, than the comparable 1982 values (.93 mg/m² vs. .54 mg/m²) although the 7m samples values were quite similar in both years (-.25 mg/m²). The elevated bulk (O-2 cm) sediment levels probably result from a mixing of surface floe oil into the upper sediment column. However, the core samples revealed only trace levels of oil lower than 5 cm in the sediment column and less oil in general in the O-5 cm section than in the O-2 cm grab sample (i.e. the benthic transect and tissue plot samples). Oil is thus probably confined to the top few (O-2) centimeters. However, general lack of strong agreement between GC²-determined (cores) and UV/F-determined (sediment samples) oil values may very well explain this difference.

The continuing use of UV/F as the main analytical tool to determine oil concentrations has created several interpretational dilemmas. Where high oil concentrations are present (e.g., Bay 11 sediments and floe) the UV/F and GC² methods agree generally within a factor of two. At lower concentrations, direct calculation of oil levels by GC² is impossible due to the biogenic interferences. The continuing use of the phytane:oil ratio to convert observed phytane levels in sediments to "oil concentrations" is risky at low levels due to the fact that phytane is not inert and the inevitability of its degradation causes the phytane to oil conversion to be grossly inaccurate, especially at low levels. Additionally, the UV/F 350-360 nm intensity measurement on total extracts could include polar aromatic compounds (metabolizes of, or polar components of petroleum) and naphtheno-aromatics which would appear as unresolved material in the f2 (aromatic-olefinic) fraction. Both these compound groupings which would be part of a "total oil" measurement are not detected in our conventional saturated and aromatic hydrocarbon determinations.

This situation is readily apparent in the tissue analyses where UV/F "oil values" on total extracts don't always agree with GC² value (phytane conversions), and don't agree with the UV/F analysis of the f2 fraction. We had previously noted (Boehm, 1983a) the widening discrepancy between UV/F and GC² values. The discrepancy between UV/F (total extract) and UV/F (f2 fraction) determinations is new data. It may have very well existed in previous years.

All signs point to the continued use of UV/F data to compare sediment and tissue data between sampling periods as being the wisest, most consistent path to take.

Evidence for petrogenic inputs to tissue samples by GC2 and GC²/MS has become difficult to establish. High, unambiguous, UV/F-determined oil values did not result, in 1983, in high aromatic hydrocarbon values by GC²/MS. The UV/F-determined values for the five species indicated the following:

1. Bay 11: Mya oil concentrations were only slightly higher than they were in 1982; Macoma values were the same as in 1982 indicating a steady state uptake/deposition situation for these deposit-feeders; Serripes levels were the same in 1983 as in 1982 indicating also a steady state situation. Astarte values decreased by a factor of approximately two. Urchin levels increased somewhat (100 versus 67) over 1982. However, between August and September of 1982 the urchin values were seen to be on the increase.
2. Bay 9: Macoma, Astarte and Serripes all show concentration decreases between 1982 and 1983 in spite of apparent increases in the sedimentary oil concentrations in this Bay. Only Mya (slight increase) and urchins (50 versus 150 ppm) showed increases in the average oil levels in this Bay.
3. Bay 7: Serripes, Mya and Astarte all illustrated no increase of oil while urchins (slight increase) and Macoma samples (1.9 versus 5.2) were seen to increase in concentration.

The overall conclusion reached when reviewing these data is that the benthic detrital feeders in Bay 11 appear to remain significantly impacted by oil while for the most part the filter-feeders continue to decrease their oil burdens in this bay. The deposited oil seems generally unavailable to the filter-feeding animals. However, background values have not been reached in all of the Bay 11 species and several of the Bay 9 and 7 species, namely the urchins. It should be noted that we now consider that the initial (2-week postspill) increases in the Bay 11 benthic animal oil content were almost certainly due to intrusion of Bay 9 dispersed oil, not due to the Bay 11 untreated oil itself, waterborne oil levels due to the dispersed oil release in Ragged Channel in general were shown to be ~50 ppb and were higher, up to 140 ppb in Bay 11. Therefore, the impact of the Bay 11 release is probably just now (1983) being revealed in the increasing or steady-state oil levels seen in the Bay 11 benthic animals (See Figure 3.53).

While aromatic hydrocarbons were detected at low to moderate levels in many of the tissue samples, levels of the petroleum aromatics (i.e. **alkylated** dibenzothiophenes and **phenanthrenes**) are much lower than was observed in 1982. The aromatic hydrocarbon **levels** have apparently decreased markedly in spite of the "oil signal" being detected in the UV/F analysis (see discussion above). The vigorous in vivo biodegradation of oil components in all of the animals discussed at length in Boehm et al.(1982a,b) and Boehm, (1983) and recently demonstrated in our analyses of the animals from the DIAND tank experiments (Boehm, 1984) may very **well** be responsible for aromatic hydrocarbon degradation as well, after the saturated hydrocarbon substrate has been depleted. The lack of large amounts of GC2/MS-determined aromatic hydrocarbons also does not mean that other aromatic compounds, (not original analytical targets of the GC²/MS) are not present.

Both the Milne Inlet sediment and animal samples were free of any UV/F-determined oil, thus lending credence to the UV/F-measured oil levels in the test bays.

Large quantities of oil still remains on the Bay 11 beach. Based on this year's findings and/or by extrapolation of the relative impact curves shown in Figure 5.1, it can be predicted that:

1. Oil will continue to "weather" on the Bay 11 beach.
2. Oil will continue to be transported offshore to the Bay 11 sediments and that levels will further increase with time.
3. Oil may will be transported from Bay 11 subtidal sediments further offshore into the deeper parts of the Bay 11/12 system.
4. Deposit feeders in Bay 11 will continue to be impacted by the oil. Oil will become less available to filter-feeders as the primary transport mechanisms of weathered oil will be through the surface sediment.
5. In vivo degradation of saturated and aromatic hydrocarbons will be the main detoxification mechanism available to animals.
6. Low levels of water-borne oil will continue to leach off the Bay 11 beach and will result in low level petroleum contamination of the water column.




FIGURE 5.1. SUMMARY OF COMPARATIVE FATES OF OIL FROM THE BIOS SPILLS.

The spill study should enter a new phase. The comparative aspects of the fate of dispersed versus untreated oils is at an end. The long term impacts of chronic oil **pollutant inputs to** an arctic nearshore environment has just begun. It is extremely important that this unique research opportunity not be lost as have others (e.g., Amoco Cadiz) in the past.

SECTION SIX

REFERENCES

- Boehm, P.D. 1984. Analysis of hydrocarbon content and composition of biological tissue and sediment samples from the 1983 **DIAND/BIOS** Tank Study, Final Report submitted to **DIAND**, Ottawa.
- Boehm, P.D. 1983. Chemistry 2: Analytical **Biogeochemistry** - 1982 Study Results, **(BIOS) Baffin Island Oil Spill** Working Report 82-2, 210 pp. Environmental Protection Service, Edmonton, Alberta.
- Boehm, P.D. 1983. The long term fate of crude oil spilled in the Arctic nearshore environment, pp. 280-291 In Proceedings Sixth Arctic Marine Oil Spill Program Technical Seminar, Environmental Protection Service, Ottawa, Canada.
- Boehm, P.D. 1981. Chemistry 2: Hydrocarbon Chemistry - 1980 Study Results. **(BIOS) Baffin Island Oil Spill** Working Report 80-2, 184 pp. Environmental Protection Service, Edmonton, Alberta.
- Boehm, P. D., J.E. Barak, D.L. Fiest, and A. Elskus. 1982c. A chemical investigation of the transport and fate of petroleum hydrocarbons in littoral and benthic environments: The Tsesis oil spill. Mar. Environ. Res. 6:157-188.
- Boehm, P. D., D.L. Fiest, and A. Elskus. 1981. Comparative weathering patterns of hydrocarbons from the Amoco Cadiz oil spill observed at a variety of coastal environments. In Proceedings of the International Symposium on the Amoco Cadiz: Fates and Effects of the Oil Spill, Centre National pour L'exploitation des oceans, Paris, France.
- Boehm, P. D., D. Fiest, and P. Hirtzer. 1982a. Chemistry 2: Analytical Biogeochemistry - 1981 Study Results **(BIOS) Baffin Island Oil Spill** Working Report 81-2:354 pp. Environmental Protection Service, Edmonton, Alberta.
- Boehm, P. D., D.L. Fiest, P. Hirtzer, L. Scott, R. Nordstrom and R. Engelhardt. 1982b. A biogeochemical assessment of the BIOS experimental spills: Transport pathways and fates of petroleum in benthic animals. pp. 581-618. In Proceedings of the Fifth Arctic Marine Oil Spill Program Technical Seminar, Environmental Protection Service, Ottawa, Canada.
- Brown, D. W., L.S. Ramos, A.3. Friedman, and W.D. MacLeod. 1979. Analysis of trace levels of petroleum hydrocarbons in marine sediments using a solvent/slurry extraction procedure. Pages 161-167 In Trace Organics Analysis: A new frontier in analytical chemistry. National Bureau of Standards Special Publication 519.
- Gordon, D. C., and P.D. Keizer. 1974. Estimation of petroleum hydrocarbons in seawater by fluorescence spectroscopy: improved sampling and analytical methods. Fish. Mar. Serv. Res. Dev. Tech. Rep. No. 481, Bedford Institute of oceanography.

- Humphrey, B. 1984. **Baffin** Island Oil Spill Project Chemistry Component, Draft Report on the 1983 oil spill experiments. Volume 1. Summary of Field Work and **Shorline** Hydrocarbon Analysis, Environment Canada, EPS, Edmonton, Alberta 62 pp.
- Humphrey, B. 1983. Chemistry 1: Field Sampling and Measurement - 1982 Study Results, (BIOS) **Baffin** Island Oil spill Working Report 82-1, 64 pp. Environmental Protection Service, Edmonton, Alberta.
- LLOYD, J.B.F. 1971. The nature and evidential value of the luminescence of automobile engine oils and related materials. 3. Forensic Sci. Soc. **11:83-94,153-10**, 235-253.
- Owens, E. H., Harper, J.R. and Foget, C.R. 1983. Shoreline Countermeasures - 1982 Study Results, (BIOS) **Baffin** Island Oil Spill Working Report 82-4, 129 pp. Environmental Protection Service, Edmonton, Alberta.
- Requejo A. and J.G. Quinn. 1983. Geochemistry of C₂₅ and C₃₀ biogenic alkenes in sediments of the Narragansett Bay estuary. Geochim. Cosmochim. Acts **47:1075-1090**.
- Wakeham, S.G. 1977. Synchronous fluorescence spectroscopy and its application to indigenous and petroleum-derived hydrocarbons in **lacustrine** sediments. Environ. Sci. Technol. **11:272-276**.
- Warner, S.S. 1976. Determination of aliphatic and aromatic hydrocarbons in marine organisms. Anal. Chem. **48:578-583**.